

**Efficacy of Supplementation of Exogenous Phytase Enzyme  
on Productivity and Carcass Characteristics of Different  
Strain of Commercial Broilers**

**A Thesis**

**By**

**ELARA PARVEEN**

Registration No. 1205103

Semester: July-December, 2014

Session: 2012-2013

MASTER OF SCIENCE (M.S.)

IN

POULTRY SCIENCE



**DEPARTMENT OF DAIRY AND POULTRY SCIENCE  
HAJEE MOHAMMAD DANESH SCIENCE AND TECHNOLOGY UNIVERSITY  
DINAJPUR-5200**

**DECEMBER, 2014**

**Efficacy of Supplementation of Exogenous Phytase Enzyme  
on Productivity and Carcass Characteristics of Different  
Strain of Commercial Broilers**

**A Thesis**

**By**

**ELARA PARVEEN**

Registration No. 1205103

Semester: July-December, 2014

Session: 2012-2013

Submitted to  
The Department of Dairy and Poultry Science,  
Hajee Mohammad Danesh Science and Technology University, Dinajpur  
in partial fulfillment of the requirements for the degree of

**MASTER OF SCIENCE (M.S.)**

**IN**

**POULTRY SCIENCE**

**DEPARTMENT OF DAIRY AND POULTRY SCIENCE  
HAJEE MOHAMMAD DANESH SCIENCE AND TECHNOLOGY  
UNIVERSITY, DINAJPUR-5200**

**DECEMBER, 2014**

**Efficacy of Supplementation of Exogenous Phytase Enzyme  
on Productivity and Carcass Characteristics of Different  
Strain of Commercial Broilers**

**A Thesis**

**By**

**ELARA PARVEEN**

Registration No. 1205103

Semester: July-December, 2014

Session: 2012-2013

*Approved as to style and content by:*

.....  
**(Dr. Mst. Afroza Khatun)**

Professor  
Research Supervisor

.....  
**(Dr. Tahera Yeasmin)**

Associate Professor  
Research Co-supervisor

.....  
(Prof. Dr. Mst. Afroza Khatun)

Chairman  
Examination Committee  
and  
Chairman, Department of Dairy and Poultry Science

**HAJEE MOHAMMAD DANESH SCIENCE AND TECHNOLOGY UNIVERSITY  
DINAJPUR-5200**

**DECEMBER, 2014**

***DEDICATED***  
***TO MY***  
***BELOVED PARENTS***

## **ACKNOWLEDGEMENTS**

*All praises are to the Almighty “Allah” the supreme authority of the universe, who kindly enables the author to complete the present research work successfully and to submit the thesis leading to Master of Science (M.S) degree in Poultry Science, Hajee Mohammad Danesh Science and Technology University, Dinajpur.*

*The author expresses the deepest sense of gratitude and heartfelt indebtedness to his honorable and respected teacher and research supervisor, **Prof. Dr. Mst. Afroza Khatun**, Chariman, Department of Dairy and Poultry Science, Faculty of Veterinary and Animal Sciences, Hajee Mohammad Danesh Science and Technology University, Dinajpur for her scholastic guidance, cordial consistence, sympathetic supervision, valuable advice and radical investigation in all phases of research work and preparation of this thesis.*

*The author is ever grateful and immensely indebted to his respected teacher and research co-supervisor, **Dr. Tahera Yeasmin**, Associate Professor, Department of Dairy and Poultry science, Faculty of Veterinary and Animal Sciences, Hajee Mohammad Danesh Science and Technology University, Dinajpur for her sympathetic supervision, valuable advice and encouragement during the entire study period.*

*The author would like to acknowledge his gratitude and sincere appreciation to respected teacher **Prof. Md. Saifur Rahman**, Department of Statistics, Hajee Mohammad Danesh Science and Technology University, Dinajpur for his encouragement, constructive and informative suggestions in course of the research work period.*

*The author also would like to acknowledge thanks and appreciation to Dr. Md. Shoriful Islam, Dr. Biswjit Ghosh, Dr. Md. Saiful Islam, Dr. Al Helal Mondol, Dr. Abdul Momin, Tapan, Saifur, Gobinda, Fojlu for their cordial help and support in completion this manuscript.*

*Finally, the author acknowledges his deepest sense of gratitude and heartfelt thanks to his beloved parents who sweated much to bring him up today’s position; brothers, and relatives for their wishes and great sacrifices during the entire study period.*

*The author.*

## ABSTRACT

An experiment was conducted to investigate the effect the efficacy of supplementation of exogenous phytase enzyme on productivity and carcass characteristics of different strain of commercial broilers at the open sided poultry shed in Hajee Mohammad Danesh Science and Technology University, Dinajpur. There were four strains of commercial broiler such as Fast feather, Arbor acres, Cobb-500 and Hubbard classic each having 78 number used for the experiment for a period of 5 weeks. A total number of 312 day old straight run broiler chicks were distributed to two dietary treatments i.e. basal diet (Control-T<sub>0</sub>) and basal diet supplemented with phytase enzyme @ 1gm/kg feed. The results indicated that broilers given diets supplemented with phytase have enhanced body weight and weight gain when compared with these fed basal diet (P<0.05). The final body weight was increased significantly (P<0.01) on T<sub>1</sub>C (Cobb-500 fed diet with 1g phytase enzyme/kg feed) and T<sub>1</sub>F (Fast feather fed diet with 1g phytase enzyme/kg feed) compared to control. There were significant difference (P>0.05) among different treatments in relation to feed consumption. Significant differences (P<0.01) were found in feed conversion ratios among birds fed on diet treated with phytase enzyme. Feed conversion ratios during the 5<sup>th</sup> week of age was 1.88, 1.87, 1.82, 1.86, 1.77, 1.72, 1.75 and 1.70 in T<sub>0</sub>F, T<sub>0</sub>A, T<sub>0</sub>C, T<sub>0</sub>H, T<sub>1</sub>F, T<sub>1</sub>A, T<sub>1</sub>C and T<sub>1</sub>H treatment groups respectively. Livability was similar in different treatments. Phytase supplementation had no significant effect on carcass cuts and dressing percent compared to non-phytase group. Dressing parameters were almost similar in different treatments and the differences were insignificant among treatment but the dressing weight percentage, thigh weight percentage and drumstick weight percentage were significant (P>0.01). Profitable ratios of the phytase groups were always higher than the control group. The cost of production was the highest in treatment T<sub>1</sub>F followed by treatment T<sub>1</sub>C, T<sub>0</sub>F, T<sub>1</sub>A, T<sub>0</sub>C, T<sub>0</sub>A, T<sub>1</sub>H, and T<sub>0</sub>H. Net profit per live broiler was the highest in treatment T<sub>1</sub>A followed by treatment T<sub>1</sub>C, T<sub>1</sub>F, T<sub>1</sub>H, T<sub>0</sub>C, T<sub>0</sub>F, T<sub>0</sub>A, and T<sub>0</sub>H respectively. Result of the present study suggests that the addition of dietary phytase was found to increase production performance and reduced cost of production.

**Keywords:** Phytase, different strain, broiler performance.

# CONTENTS

| Chapter           | Title  | Page        |
|-------------------|--|-------------|
|                   | ACKNOWLEDGEMENTS   | v           |
|                   | ABSTRACT   | vi          |
|                   | CONTENTS   | vii-xii     |
|                   | LIST OF TABLES   | x           |
|                   | LIST OF FIGURES  | xi          |
|                   | LIST OF ABBREVIATIONS  | xii         |
| <b>CHAPTER I</b>  | <b>INTRODUCTION</b>  | <b>1-3</b>  |
| <b>CHAPTER II</b> | <b>REVIEW OF LITERATURE</b>                                      | <b>4-10</b> |
| 2.1               | Phytase  | 4           |
| 2.2               | Phytate  | 4-5         |
| 2.3               | Structure of phytate   | 5           |
| 2.4               | Commercial phytase enzymes                                       | 5-6         |
| 2.5               | Effect of phytase on body weight                                 | 6           |
| 2.6               | Effect of microbial phytase on the growth performance of broiler | 7           |
| 2.7               | Effect of phytase on feed intake                                 | 7           |
| 2.8               | Effect of phytase on feed conversion ratio                       | 7           |
| 2.9               | Effect of phytase on livability                                  | 8           |
| 2.10              | Effect of phytase on feed cost                                   | 8           |
| 2.11              | Effect of phytase on meat yield characteristics                  | 8           |
| 2.12              | Effect of phytase on digestibility                               | 8-9         |
| 2.13              | Effect of phytase on environment                                 | 9           |
| 2.14              | Effect of phytase on energy utilization                          | 9-10        |
| 2.15              | Effect of phytase on mineral retention                           | 10          |

## CONTENTS (Contd.)

| Chapter            | Title                                 | Page         |
|--------------------|---------------------------------------|--------------|
| <b>CHAPTER III</b> | <b>MATERIALS AND METHODS</b>          | <b>11-18</b> |
| 3.1                | Statement of the experiment           | 11           |
| 3.2                | Experimental birds                    | 11           |
| 3.3                | Layout of the experiment              | 11-12        |
| 3.4                | Preparation of the experimental house | 12           |
| 3.5                | Experimental ration                   | 13-14        |
| 3.5.1              | Collection of enzymes                 | 13           |
| 3.5.2              | Experimental diet                     | 13-14        |
| 3.6                | Management of the experimental birds  | 15           |
| 3.6.1              | Litter management                     | 15           |
| 3.6.2              | Brooding                              | 15           |
| 3.6.3              | Lighting                              | 16           |
| 3.6.4              | Floor, feeder, waterer space          | 16           |
| 3.6.5              | Feed and water management             | 16           |
| 3.6.6              | Immunization                          | 16-17        |
| 3.6.7              | Medication                            | 17           |
| 3.6.8              | Sanitation                            | 17           |
| 3.7                | Post-mortem examination               | 17           |
| 3.8                | Data collection and record keeping    | 18           |
| 3.9                | Statistical analysis                  | 18           |

---



## **CONTENTS (Contd.)**

| <b>Chapter</b>    | <b>Title</b>  | <b>Page</b>  |
|-------------------|---|--------------|
| <b>CHAPTER IV</b> | <b>RESULTS AND DISCUSSION</b>   | <b>19-30</b> |
|                   | 4.1 Effect of Phytase enzyme on body weight gain of broilers                  | 19           |
|                   | 4.2 Effect of Phytase enzyme on feed intake of broilers                       | 22           |
|                   | 4.3 Effect of Phytase enzyme on feed conversion ratio (FCR) of broilers       | 22-23        |
|                   | 4.4 Effect of Phytase enzyme on carcass characteristics of broilers           | 26           |
|                   | 4.5 Effect of Phytase enzyme on production cost and profit margin of broilers | 26-27        |
|                   | 4.6 Effect of Phytase enzyme on livability of broilers                        | 30           |
| <b>CHAPTER V</b>  | <b>SUMMARY AND CONCLUSION</b>   | <b>31-32</b> |
|                   | <b>REFERENCES</b>   | <b>33-42</b> |

---

## LIST OF TABLE

| Table No. | Title   | Page |
|-----------|---|------|
| 2.1       | Production strain and gene origin of commercial enzymes   | 6    |
| 2.2       | Influence of microbial phytase on body weight in broilers   | 7    |
| 2.3       | Effects of phytase supplementation on energy utilization (AME or AMEn) in broiler chicken   | 10   |
| 2.4       | Effects of dietary levels of microbial phytase on calcium, phosphorus, & magnesium & zinc retention in broiler chicks from 0 to 3 weeks of age. | 10   |
| 3.1       | Different strains of chicken (broiler) used for the experiment.   | 11   |
| 3.2       | Layout of the experiment  | 12   |
| 3.3       | Composition of the experimental starter and finisher diets fed to broilers.   | 14   |
| 3.4       | Vaccination schedule followed for the experimental broiler strains  | 17   |
| 4.1       | Effect of Phytase enzyme in the diet of broilers on body weight (g/bird )   | 20   |
| 4.2       | Effect of Phytase enzyme on the production performance of broiler.  | 21   |
| 4.3       | Effect of Phytase enzyme in the diet of broilers on feed intake (g/bird)  | 24   |
| 4.4       | Effect of Phytase enzyme in the diet of broilers on feed conversion ratio (g/bird)  | 25   |
| 4.5       | Effect of Phytase enzyme in the diet of broilers on carcass characteristics at 35 days of age   | 28   |
| 4.6       | Effect of Phytase enzyme on production cost and profit margin of broilers.  | 29   |

---


## LIST OF FIGURE

| Figure No. | Title  | Page |
|------------|--|------|
| 1          | Preparation of the experimental diets                      | 13   |
| 2          | Brooding and housing of birds during experimental period   | 15   |
| 3          | Feeding management of birds during experimental period     | 16   |
| 4          | Postmortem examination of birds during experimental period | 17   |
| 5          | Processing of birds during experimental period             | 18   |

---

## LIST OF ABBREVIATIONS AND SYMBOLS

| ABBREVIATIONS | ELABORATIONS                    |
|---------------|---------------------------------|
| BASF          | = Badische Anilin- & Sodafabrik |
| DSM           | = Dutch State Mines             |
| FTU           | = Phytase units                 |
| IU            | = International unit            |
| FCR           | = Feed conversion ratio         |
| MS            | = Master of Science             |
| PM            | = Post mortem                   |
| AME or AMEn   | = Apparent metabolisable energy |
| mg            | = Miligram                      |
| g             | = Gram                          |
| kg            | = Kilogram                      |
| Lit           | = Liter                         |
| ml            | = Mililiter                     |
| kcal          | = kilo-calorie                  |
| CP            | = Crude protein                 |
| CF            | = Crude fiber                   |
| DM            | = Dry matter                    |
| Dr.           | = Doctor                        |
| Prof.         | = Professor                     |
| SEM           | = Standard Error of Means       |
| Tk.           | = Taka                          |
| P             | = Phosphorus                    |
| S             | = Sulphur                       |
| Ca            | = Calcium                       |
| K             | = Potassium                     |
| Mg            | = Magnesium                     |
| Mn            | = Manganese                     |
| Na            | = Sodium                        |
| NS            | = Not significant               |
| %             | = Per cent                      |
| &             | = and                           |
| @             | = At the rate of                |
| +             | = Plus/and                      |
| /             | = Per/or                        |
| >             | = Greater than                  |
| <             | = Less than                     |
| ±             | = Plus-minus                    |



# CHAPTER 1

## INTRODUCTION

# CHAPTER 1

## INTRODUCTION

In commercial poultry ration, nutrients are fortified through accumulation of different feed ingredients furnished with necessary micro nutrients. Protein costs involve about 45 percent of the total feed cost. The daily requirements of dietary protein are furnished from different animal and plant sources. But the sources of plants are sometimes become harmful for the poultry as because plant sources contain some anti nutritional factors like phytate phosphorus, trypsin inhibitors, non-starch polysaccharides (NSP), oligosaccharides and lectins (Deshpande and Cheryan, 1984; NRC, 1994) which decrease feed consumption but increases growth rate and feed utilization. Phytate phosphorus reduces the phosphorus and calcium availability in poultry. It is well documented that phosphorus is one of the basic mineral elements in all feed rations, having a greater influence on biological systems. Feeds of plant origin protein contain significant amount of this mineral; however, 50-80% of phosphorus is bound in phytates that cannot be broken down by endogenous enzymes in poultry (Deshpande and Cheryan, 1984). As a consequence, phosphorous from plant sources is poorly digested and cannot meet nutritional requirements of poultry regardless the fact that phytate phosphorus amounts in cereal grains can be as high as 50-80%, in legumes 50-68%, in oil-producing plant seeds and their by-products 51-76% (Eeckhout and Peape., 1994; Jeroch, 1993 and Oloffs, 2000). The major proportion of the phosphorus is stored in a special way: six phosphorus molecules are bound to phytic acid in a ring form. This phytic acid ring is called phytate. Besides, phytate creates a large number of insoluble salts with divalent and trivalent cations such as calcium, magnesium, potassium, Iron, manganese or zinc. The interaction of protein/phytate and starch/phytate obstructs the digestion of protein and carbohydrates (Knuckles, 1989; Zyla, 1992). Therefore, just like phosphorus, these valuable nutritive substances are also lost to animal as excreted in the faeces. Phytic acid also suppresses the activity of certain enzymes such as a amylase, trypsin, tyrosinase and pepsin, thereby suppressing crude starch and crude protein digestion (Zyla, 1992). In order to become P available to broiler chicks, Phosphorous from plant sources must be hydrolyzed, with phytase as a catalyst, to inositols and inorganic phosphates which are readily absorbed in digestive tract. Through supplementation of microbial phytase

to the monogastric animals about 50% of phytic phosphorous may be released. Results of numerous experiments have shown that degradation of phytate by phytase has two-fold positive effect-release of phosphorous and release of minerals. It is the enzyme known to release the orthophosphate group from the phytate molecule. Improving the availability of phytate, P would reduce the necessity to include feed phosphates in the diet and enable a reduction of the dietary P contents, without jeopardizing the bird's health and productivity. In turn, this would result in a lower P excretion per unit of edible product (eggs, meat), and reduce P-linked environmental pollution problems by intensive livestock production (Kornegay and Ravindran, 1996; Van, *et al.*, 1997).


The supplementation of poultry diets with exogenous enzymes to enhance their performance is not a new concept and research articles in this field date back to the early part of the 20<sup>th</sup> century (Hastings, 1946), with the first published article in 1925 (Rosen, 2010). A phytase (myo-inositol hexakisphosphate phosphohydrolase) is any type of phosphatase enzyme that catalyzes the hydrolysis of phytic acid (myo-inositol hexakisphosphate) an indigestible, organic form of phosphorus that is found in grains and oil seeds and releases a unstable form of inorganic phosphorus. Phytate ( salt form of phytic acid) and phytate bound phosphorus (P) is present in all poultry diets and the partial availability of phytate-P has long been recognized (Lowe *et al.*, 1939). Warden and schaible (1962) were the first to show that exogenous phytase enhances phytate-P utilization and bone mineralization in broiler chicks.

The positive effect of phytase can be expressed through better appetite, improve feed conversion ratio, increase carcass quality, increase digestibility. Also, adding phytase decreases the amount of total and soluble phosphorus in the litter, which has positive effect on the environment when poultry litter is used as fertilizer. Equally important, phytase may reduce the cost of the diet by reducing the amount of soybean meal, fat and crystalline amino acids that must be added. Research also has indicated that the improvement in growth performance observed in chicken, fed phytase were associated with increased feed intake and feed efficiency which might be due to release and utilization of P from the phytate mineral complex (Qian *et al.*, 1996; Sebastian *et al.*, 1996) or

utilization of inositol (Simans *et al*, 1990) or increase starch digestibility (Knukles and Betschart, 1987) or increased utilization of protein and amino acids (Rama Rao *et al*, 1999) or overall utilization of nutrients (Miles and Nelson, 1974). Phytase has been shown to increase the availability of some trace minerals, including copper, manganese, iron and zinc. Because of the positive effect of phytase on trace mineral utilization commercial use may lead to removing trace minerals in diets where phytase is added. It has been shown that removing the trace mineral premix from poultry diets from hatch to 42 days has no effect on growth performance, but it does have negative effects on bone growth. This negative effect was not overcome with the addition of phytase, indicating that phytase may not be able to replace the trace mineral premix in diets for broilers. Therefore, the present research work was designed with the following objective:

- i) To investigate the effect of exogenous phytase enzyme on different strains of commercial broilers.
- ii) To examine the productivity and carcass characteristics of different strains of commercial broiler.
- iii) To evaluate the benefit of rearing broiler after feeding phytase enzyme.





# CHAPTER 2

## REVIEW OF LITERATURE

## CHAPTER 2

### REVIEW OF LITERATURE

The review of literatures as presented covers recognition of potential sources of vital information about the analogous researches study conducted by other workers and helps to identify the basic concepts of different components blended in researches. The contributions made by numerous research workers are pertinent in this research are discussed below.

#### 2.1 Phytase

A phytase (myo-inositol hexakisphosphate phosphohydrolase) is any type of phosphatase enzyme that catalyzes the hydrolysis of phytic acid (myo-inositol hexakisphosphate)-an indigestible, organic form of phosphorus which is found in grains and oil seeds and releases a usable form of phosphorus (Mullaney *et al.*, 2000). Phytases have been found in animals, plants, fungi and bacteria (Mullaney *et al.*, 2003).

#### 2.2 Phytate

Phytate, the mixed cation salt of phytic acid (Myo-inositol-1, 2, 3, 4, 5, 6-hexakis dihydrogen phosphate), is a naturally occurring compound present in feedstuffs of plant origin (Reddy *et al.*, 1982). Phytate was identified as isolated small, unknown particles from a variety of plant seeds (Hartig, 1855; Reddy *et al.*, 1982). Phytate serves as the primary storage form of P and inositol in seeds (Hidvegi & Lasztity, 2002). It is also involved in controlling homeostasis of P levels in seeds (Lott *et al.*, 2000) and plays an important role in plant growth and seed germination (Aureli *et al.*, 2011). In some plants, phytic acids binds potassium (K<sup>2+</sup>), magnesium (Mg<sup>2+</sup>) and to a lesser extend calcium (Ca<sup>2+</sup>) to form phytin (Maenz, 2001). Phytin is stored in vacuoles known as protein bodies. It is distributed in dense aggregates called globoids or can be distributed throughout the proteinaceous matrix (Maenz, 2001). Phytate accumulates in the aleuronic layer in monocotyledonous seeds (wheat, rice, barley) and in the germ of corn (Hidvegi and Lasztity, 2002). The amount of phytate in plant sources is influenced by cultivar and climatic conditions. Phytate is located in the

outer parts of the kernel and therefore different milling methods can also influence the phytate content of the end products (Hidvegi & Lasztity, 2002). Phytate levels can be measured through the use of high performance liquid chromatography (HPLC) and the amount of phytate bound P can be calculated as 28.2% of the total phytate concentration (Sauvant *et al.*, 2004).

### **2.3 Structure of phytate**

Phytic acid is a charged molecule and consists out of a myo-inositol ring (a six carbon molecule) and six phosphate groups extending from the structure (Johnson and Tate, 1969). The molecule has 12 proton dissociation sites with a high chelation capacity for multivalent cations (Cheryan and Rackis, 1980) and positively charged nutrients (Selle and Ravindran, 2007). At neutral pH, phytic acid can have one or two negatively charged oxygen atoms in the phosphate groups. Therefore there is likely to be a strong correlation interaction between cations and two phosphate groups and also a weak correlation interaction between cations and a single phosphate group (Singh, 2008).

### **2.4 Commercial phytase enzymes**

The first successful commercial phytase (Natuphos®) was developed in 1991 by Badische Anilin- & Sodafabrik (BASF). For many years, Natuphos was the only phytase enzyme in the market (Simon and Igbasan, 2002). It is a 3-phytase produced by the fungus *Aspergillus*. In order to produce this enzyme on large scale, BASF genetically modified the recombinant *A. niger* strain with the *A. ficuum* phytase gene (Zhang *et al.*, 2000). DSM marketed Ronozyme, a 6-phytase produced by the *Aspergillus oryzae* strain, transformed with a gene from *Peniophora lycii* (Simon and Igbasan, 2002). Phyzyme is classified as a 6-phytase and is produced by *Schizosaccharomyces pombe* and modified with the App A gene from *E. coli* (Kerr *et al.*, 2010). DSM was the first company to use synthetic genes in a phytase product. Ronozyme HiPhos is a 6-phytase (histidine acid phosphatase phytase), produced by *Aspergillus oryzae* and contains two synthetic genes that mimic a phytase gene from *Citrobacter braakii* ATCC 51113 (Guggenbuhl *et al.*, 2012). All types of commercially available phytases belong to the class of histidine acid phytases (based

on the catalytic mechanism). Acid phytases show a maximum phytate dephosphorylation at pH 5, therefore histidine acid phytases are suspected to be most efficient in the crop (pH 4.0 to 5.0) or in the proventriculus and gizzard (pH 2.0 to 5.0) of the chicken (Greiner and Konietzny, 2011). The classes of certain commercial enzymes with its production strain and gene origin are summarised in Table 2.1. Currently there are quite a few phytase enzymes available commercially. The recommended dosages to release similar amounts of phytate bound P differs greatly among different phytase source.

**Table 2.1 Production strain and gene origin of commercial enzymes**

| Name of Phytase | Production strain/organism       | Origin of phytase gene        | Class     | Company | Reference  |
|-----------------|----------------------------------|-------------------------------|-----------|---------|--|
| Natuphose       | <i>Aspergillus niger</i>         | <i>Aspergillus ficuum</i>     | 3-phytase | BASF    | Zhang <i>et al.</i> , 2000                         |
| Phyzyme         | <i>Schizosaccharomyces pombe</i> | AppA gene from <i>E. coli</i> | 6-phytase | Danisco | Kerr <i>et al.</i> , 2010                          |
| Ronozyme        | <i>Aspergillus oryzae</i>        | <i>Peniophora lycii</i>       | 6-phytase | DSM     | Simon and Igbasan. 2002; Kerr <i>et al.</i> , 2010 |
| HiPhos          | <i>Aspergillus oryzae</i>        | <i>Citrobacter braakii</i>    | 6-phytase | DSM     | Guggenbuhl <i>et al.</i> , 2012                    |

## 2.5 Effect of phytase on body weight

Piao *et al.* (1998) observed that enzyme supplemented with phytase diets increased live weight of broiler. The improvement in growth performance observed in the chickens fed phytase were associated with increased feed intake and feed efficiency, which might be due to the release and utilization of P-from the phytate-mineral complex (Qian *et al.*, 1996; Sebastain *et al.*, 1996) or utilization of inositol (Simons *et al.*, 1990) or increase starch digestibility (Knukles and Betschart, 1987) or increased utilization of protein and amino acids (Rama Rao *et al.*, 1999) or over all utilization of nutrients (Miles and Nelson, 1974). Addition of exogenous phytase in low protein diet have been reported to improve growth performance of broiler (Lan *et al.*, 2002; Ahmed *et al.*, 2000). But result of this study contradicts with the report of Pizzolante *et al.*(1999),who reported that the dietary phytase had no effect on live weight gain of broilers. Richered *et al.* (1994) found that growth was not improved by enzyme supplementation in triticale based diet.

## 2.6 Effect of microbial phytase on broiler growth performance

Many authors reported that microbial phytase supplementation increased body weight gain, feed intake and feed conversion efficiency in broilers.

**Table 2.2** Influence of microbial phytase on body weight in broilers.

| Age (Days) | Phytase (g/kg) | Non-phytase (Phosphorus in diet) | Weight gain(%) | FCR  | Sources                        |
|------------|----------------|----------------------------------|----------------|------|--------------------------------|
| 28         | 750            | 0.169                            | 38             | 0.63 | Simons <i>et al.</i> (1990)    |
| 22         | 500            | -                                | 13             | 3.4  | Broz <i>et al.</i> (1994)      |
| 20         | 600            | -                                | 77             | 3.7  | Aoyagi and Baker 1995          |
| 20         | 600            | 0.45                             | 11             | 3.53 | Sebastian <i>et al.</i> (1996) |
| 21         | 600            | 0.20                             | 36             | 2.60 | Kornegay <i>et al.</i> (1996)  |

## 2.7 Effect of phytase on feed intake

Phytase supplementation increased feed consumption by partial degrading of cell wall of feed and increased digestibility of nutrients Ahmad *et al.* (2000); Naher, B. (2002); Aksakal, N. and Bilal, H. (2000). On the other hand, the result contradicts with the findings of some earlier workers (Wilson *et al.*, 1999). They found that feed consumption was decreased due to addition of enzymes since birds fulfilled their nutrient requirement by taking less amount of feed. Phytase supplementation increased feed intake of broilers fed on phosphorus deficient diets (Mondal *et al.*, 2007) where the minimum dietary level of phytase was 1000 PU/kg. Some investigators reported similar feed intake on different levels of phytase observed in duck, turkey and chicken respectively (Atia *et al.*, 2000; Orban *et al.*, 1999 and Ciftci *et al.*, 2005).

## 2.8 Effect of phytase on feed conversion ratio

It is reported that addition phytase at a level higher than 500 PU/kg had impact in feed conversion efficiency (Mondal *et al.*, 2007). Lan *et al.* (2002); Moshad (2001); Aksakal and Bilal (2002); Scott *et al.*, (1997) reported that feed conversion was increased due to better feed utilization by birds

## **2.9 Effect of phytase on livability**

Pillai *et al.* (1995) found survivability was similar in control and phytase enzyme treated feed. Alam (2001) found similar livability among the different dietary groups.

## **2.10 Effect of phytase on feed cost**

Augelovicova and Michalik (1997) Stated that enzyme supplementation in commercial broiler diet decreased feed cost by 8.81% to 9.73% for production of 1 kg broiler meat. Nahar (2002) evident that feed cost per kg was reduced for addition of enzyme and increased profitability of duck rearing. Ahmed *et al.* (2000); Kies *et al.* (2001); Ren *et al.* (1999); Augelovicova and Michalik (1997) and Morkunas *et al.* (1993) reported that reduced feed cost was for the improved feed utilization and faster growth rate of broilers.

## **2.11 Effect of phytase on meat yield characteristics**

A increase of total meat and breast meat for supplementing enzyme have been supported by Rahman *et al.* (2009). Khawaja (2003) and Rabayaa (2003) noted higher dressing yield of broilers for dietary supplementation of phytase. Pillai *et al.*, (2006) and Angel *et al.* (2007) contradict those findings. They did not find any change in meat yield in broilers which could be attributed to phytase supplementation. A positive correlation of dressed weight with live weight or age obtained coincides with the findings of some earlier workers (McNally and Spicknall, 1949; Jaap *et al.*, 1950; Howlider and Rose, 1989)

## **2.12 Effect of phytase on digestibility**

Phytate-Ca forms insoluble complex with starch and fatty acids in the gastrointestinal tract of broilers, which depress the digestibility of carbohydrate and lipid (Rama Rao *et al.*, 2001). Therefore supplemental phytase had positive effects on dry matter digestibility by releasing bound organic nutrients (Ravindran *et al.*, 2000). Improved nitrogen digestibility due to added phytase has been reported in several studies (Kornegay, 1996; Sebastian *et al.*, 1997; Ravindran *et al.*, 2000). The interaction of

phytic acid and protein forms phytate-protein (Cheryan, 1980). They phytate also binds with major proteolytic enzyme, trypsin (Caldwell, 1992) eventually leading to lowered digestibility's of nitrogen and amino acids. It is therefore, lively that when phytase hydrolyses the ester bonds to release P from the phytic acid molecule, it will also release the phytate bond protein and removes the negative effects of phytic acid on proteolytic enzymes, thus increasing the digestion and absorption of protein and amino acids (Ravindran *et al.*, 2000).

### **2.13 Effect of phytase on environment**

With the public concern over pollution due to P excretion with feces considerable attention has been generated in the recent years on broiler and layer chickens (Sebastian *et al.*, 1996; Zhang *et al.*, 1999; Ravindran *et al.*, 2000). Phytase supplementation decreased P excretion in the manure and reduced the potential environmental problems (Jalal *et al.*, 2001; Biehl and Baker 1997). Exogenous phytase supplementation seems to be a solution to combat phytate issues, thus leads to bioavailability of P economically and reduces P load on environment (Shelton *et al.*, 2004).

### **2.14 Effect of phytase on energy utilization**

Early studies involving dephytinised feed ingredients showed that phytate negatively influences energy utilization in broilers (Rojas and Scott, 1969; Miles and Nelson, 1974). Exogenous phytase has consistently increased AME of broiler diets based on wheat and/or sorghum in many studies. (Ravindran *et al.*, 1999, 2000, 2001; Selle *et al.*, 1999, 2001, 2005). These studies *and* several other researches (Driver *et al.*, 2006; Namkung and Leeson, 1999; Shirley and Edwards, 2003) are summarized in Table 2.3. Overall, phytase supplementation increased AME by an average of 0.36 MJ kg<sup>-1</sup> DM (or 2.8%) over the non-supplemented controls. The percentage responses in AME following phytase supplementation are negatively correlated to the energy density of the control diets.

**Table 2.3 Effects of phytase supplementation on energy utilization (AME or AMEn) in broiler chicken.**

| No. | Diet type              | AME (MJ kg <sup>-1</sup> DM) |         | Response               |     | Phytase (FTU kg <sup>-1</sup> )  |
|-----|------------------------|------------------------------|---------|------------------------|-----|----------------------------------|
|     |                        | Control                      | Phytase | MJ kg <sup>-1</sup> DM | %   |                                  |
| 1   | Maize-soy              | 12.49                        | 12.62   | 0.13                   | 1.0 | 24,000 <i>Aspergillus niger</i>  |
| 2   | Sorghum                | 12.80                        | 13.10   | 0.3                    | 2.3 | 750, <i>Aspergillus niger</i>    |
| 3   | Wheat                  | 14.88                        | 14.96   | 0.08                   | 0.5 | Mean of two phytase              |
| 4   | Wheat (pre-pelleted)   | 14.2                         | 14.1    | -0.1                   | 0.7 | 600, <i>Aspergillus niger</i>    |
| 5   | Wheat-sorghum 2.3 g kg | 13.33                        | 13.52   | 0.19                   | 1.4 | 400+800 <i>Aspergillus niger</i> |
| 6   | Wheat-sorghum 4.5 g kg | 12.67                        | 13.38   | 0.71                   | 4.6 | 400+800 <i>Aspergillus niger</i> |
| 7   | Wheat-sorghum blend    | 14.22                        | 14.55   | 0.33                   | 2.3 | 500, <i>Aspergillus niger</i>    |
| 8   | Barley per se          | 12.36                        | 12.69   | 0.33                   | 2.7 | 700, <i>Aspergillus niger</i>    |

Derived from the studies Ravindran *et al.* (2000: 2001)

## 2.15 Effect of phytase on mineral retention


The microbial phytase supplementation of broiler diets has very significant beneficial effects on the mineral retention. A study showed that with increasing level of supplemental phytase increase the Ca (up to 9%), P (up to 10%) and Zn (up to 16%) retention (Brenes *et al.*, 2003). This effect is summarized in Table 4.4.

**Table 2.4 Effects of dietary levels of microbial phytase on calcium, phosphorus, & magnesium & zinc retention in broiler chicks from 0 to 3 weeks of age.**

(Adopted from Brenes *et al.*, 2003)

| Mineral    | PHY (U/kg) | Plasma Level (mg/dl) |
|------------|------------|----------------------|
| Calcium    | 0          | 0.64                 |
|            | 200        | 0.66                 |
|            | 400        | 0.67                 |
|            | 600        | 0.70                 |
| Phosphorus | 0          | 0.61                 |
|            | 200        | 0.63                 |
|            | 400        | 0.65                 |
|            | 600        | 0.66                 |
| Magnesium  | 0          | 0.37                 |
|            | 200        | 0.37                 |
|            | 400        | 0.39                 |
|            | 600        | 0.39                 |
| Zinc       | 0          | 0.24                 |
|            | 200        | 0.25                 |
|            | 400        | 0.28                 |
|            | 600        | 0.28                 |





**CHAPTER 3**  
**MATERIALS AND METHODS**

## **CHAPTER 3**

### **MATERIALS AND METHODS**

#### **3.1 Statement of the experiment**

The experiment was conducted at the open sided poultry shed under the Dept. of Dairy and Poultry Science in Hajee Mohammad Danesh Science and Technology University, Dinajpur. There were four strains of commercial broiler such as Fast feather, Arbor acres, Cobb-500 and Hubbard classic each having 78 number used for the experiment for a period of 5 weeks. A total number of 312 day-old straight run broiler chicks were used to find out the effect of exogenous phytase enzyme on the production performance of broiler.

#### **3.2 Experimental birds**

Three hundred twelve day-old chicks were collected via local traders for the experiment.

**Table 3.1 Different strains of chicken (broiler) used for the experiment.**

| <b>Strains of chicken (Broiler)</b> | <b>No. of chicks</b> | <b>Name of the farms</b>   |
|-------------------------------------|----------------------|----------------------------|
| Fast feather                        | 78                   | Paragon Poultry Ltd.       |
| Arbor acres                         | 78                   | Kazi Farm Ltd.             |
| Cobb-500                            | 78                   | Kazi Farm Ltd.             |
| Hubbard classic                     | 78                   | Aftab Bahumukhi Farms Ltd. |

#### **3.3 Layout of the experiment**

The chicks were randomly distributed to eight dietary treatment groups (T<sub>0</sub>F, T<sub>0</sub>A, T<sub>0</sub>C, T<sub>0</sub>H, T<sub>1</sub>F, T<sub>1</sub>A, T<sub>1</sub>C, T<sub>1</sub>H) having three replications in each treatment. The chicks were reared in separated pens according to treatments and replications, each dietary treatment group consisted of 39 chicks and numbers of chicks in each replication was 13.

The layout of the experiment is shown in the following Table 3.2

**Table 3.2 Layout of the experiment**

| Dietary Treatment | No of chicks in each replication |                |                | Total number of chicks per treatment |
|-------------------|----------------------------------|----------------|----------------|--------------------------------------|
|                   | R <sub>1</sub>                   | R <sub>2</sub> | R <sub>3</sub> |                                      |
| T <sub>0</sub> F  | 13                               | 13             | 13             | 39                                   |
| T <sub>0</sub> A  | 13                               | 13             | 13             | 39                                   |
| T <sub>0</sub> C  | 13                               | 13             | 13             | 39                                   |
| T <sub>0</sub> H  | 13                               | 13             | 13             | 39                                   |
| T <sub>1</sub> F  | 13                               | 13             | 13             | 39                                   |
| T <sub>1</sub> A  | 13                               | 13             | 13             | 39                                   |
| T <sub>1</sub> C  | 13                               | 13             | 13             | 39                                   |
| T <sub>1</sub> H  | 13                               | 13             | 13             | 39                                   |

**Here,**

- T<sub>0</sub>F : Fast feather + giving diet without phytase enzyme.
- T<sub>0</sub>A : Arbor acres + giving diet without phytase enzyme.
- T<sub>0</sub>C : Coob-500 + giving diet without phytase enzyme.
- T<sub>0</sub>H : Hubbard classic + giving diet without phytase enzyme.
- T<sub>1</sub>F : Fast feather + giving diet having 1g phytase enzyme/kg feed.
- T<sub>1</sub>A : Arbor acres + giving diet having 1g phytase enzyme/kg feed.
- T<sub>1</sub>C : Cobb-500 + giving diet having 1g phytase enzyme/kg feed.
- T<sub>1</sub>H : Hubbard classic + giving diet having 1g phytase enzyme/kg feed.

### **3.4 Preparation of the experimental house**

The experimental house was properly washed and cleaned using tap water. Ceiling, walls and floor were thoroughly cleaned and subsequently disinfected with Timsen (1gm/lit). After proper drying, the house divided into 24 pens of equal size using wire net. Each of pens was 12 sq feet floor spaces for 13 chicks. Fresh dried rice husk was used as litter material on the floor at a depth of 4 inch.

### 3.5 Experimental ration

#### 3.5.1 Collection of enzyme

The phytase enzyme (SQZYME SSF) was manufactured by F. Hoffmann-La Roche Ltd, Switzerland and were supplied by Square limited, Bangladesh for conducting the research work.

#### 3.5.2 Experimental diet

Broiler starter and finisher diets were provided between 0 to 14 days and 15 to 35 days of age respectively. The diets were formulated using locally available feed ingredients such as Maize, Rice polish (Auto), Soybean meal (44), Protein Concentrate, Oyster shell, DCP, Soybean oil, Common salt, Vitamin-mineral premix (Rena premise) and coccidiostat (DOTT) were added to the experimental diets as per recommendation of the manufacturer and mixed properly with the whole ration. It was done by mixing premix and DOTT (Coccidiostat), first with a small quantity of mixed feed and then with gradually increased in quantity by adding remaining mixed feed. Finally, total amount of feed was mixed thoroughly. The whole procedure was followed for both starter and finisher diets. The amount of ingredients used and the detail composition of different experimental ration are shown in Table 3.3.



Fig 1: Preparation of the experimental diets

**Table 3.3** Composition of the experimental starter and finished diets fed to broilers.

| Feed ingredients          | Amount (kg/100kg feed) |                       |
|---------------------------|------------------------|-----------------------|
|                           | Starter (0-14 days)    | Finisher (15-35 days) |
| Maize                     | 53.5                   | 57.00                 |
| Rice polish (Auto)        | 10.0                   | 10.0                  |
| Soybean meal (44)         | 23.0                   | 18.0                  |
| Protein concentrate       | 10.0                   | 10.0                  |
| Oyster shell              | 1.0                    | 0.75                  |
| DCP                       | 0.5                    | 0.75                  |
| Soybean oil               | 1.5                    | 3.0                   |
| Common salt               | 0.25                   | 0.25                  |
| **Vitamin- mineral premix | 0.25                   | 0.25                  |
| <b>Total</b>              | <b>100kg</b>           | <b>100kg</b>          |
| Nutrients                 | Calculated composition |                       |
|                           | Starter (0-14 days)    | Finisher (15-35 days) |
| ME (kcal/kg)              | 2977                   | 3074                  |
| CP (%)                    | 21.21                  | 19.40                 |
| CF (%)                    | 5                      | 5                     |
| Ca (%)                    | 1.00                   | 0.95                  |
| Available P (%)           | 0.74                   | 0.75                  |
| Ash (%)                   | 6                      | 6                     |
| Lysine (%)                | 1.02                   | 0.89                  |
| Methionine (%)            | 0.35                   | 0.35                  |

**Note:** Phytase enzyme used in enzyme grouped diet @ 10g/100kg.

\*\*Vitamin-mineral premix composition (each 2.5 kg contained): Vitamin A 12000000 IU, Vitamin D<sub>3</sub> 2000000 IU, Vitamin E 15000mg, Vitamin B<sub>1</sub> 100mg, Vitamin B<sub>2</sub> 4000mg, Vitamin B<sub>6</sub> 3000 mg, Vitamin B<sub>12</sub> 10mg, Vitamin K<sub>3</sub> 201K) mg, Folic acid 1500mg, Nicotinic acid 25000mg, Pantothenic acid 11000mg, Biotin 15mg, Iron 32000mg, Copper 8000mg, Manganese 64000mg, Cobalt 300mg, Zinc 40000mg, Iodine 800mg, Selenium 200mg, Lysine 30000mg, Methionine 50000mg, Antioxidant 10000mg.

### **3.6 Management of the experimental birds**

At the beginning of the experiment, chicks were individually weighted and recorded as initial body weight. The following management procedures were followed during the experimental period.

#### **3.6.1 Litter management**

The birds were reared on rice husk litter floor having a depth of 4 inch. At the end of each week, litter was stirred to break its compactness and maintain proper moisture. At the end of the 4 weeks of age, droppings were cleaned from the surface of the litter.

#### **3.6.2 Brooding**

Additional heat was provided for brooding the chicks when it was necessary up to the end of 1<sup>st</sup> week. Brooding temperature was maintained as bird's requirement from 33°C to normal environmental temperature of the house. Additional heat was managed by fitting 100 watt electric bulbs at the center of the pen, about 6 inches above of the floor. The height of the bulbs increased by raising the bulb gradually as per temperature requirement.



Fig 2: Brooding and housing of birds during experimental period

### **3.6.3 Lighting**

The birds were exposed to a continuous lighting of 23 hours and 30 minutes and a dark period of 30 minutes in 24 hours. Supplementary light at night was provided using electric bulb by hanging at a height of 2.8 meters to provide necessary lighting.

### **3.6.4 Floor, feeder, waterer space**

Each pen allotted for 13 birds, therefore floor space for each pen was 12 sq ft. One round feeder and one round waterer provided in each pen for 13 birds, additional feeder and waterer added according to size of the flock.

### **3.6.5 Feed and water management**

The birds were reared in separate pens. Feed was given *ad-libitum* throughout the experimental period. Phytase enzyme was given in feed.




Fig 3: Feeding management of birds during experimental period

### **5.6.6 Vaccination Schedule**

The experimental birds were vaccinated against Baby chick Ranikhet Disease and Gumboro Disease as per following schedule:

**Table: 3.4** Vaccination schedule followed for the experimental broiler strains.

| Name of the vaccine | Age of the bird      | Dose and route of administration of diluted vaccine |
|---------------------|----------------------|---|
| BCRDV               | 4 <sup>th</sup> day  | 1 drop in each eye                                  |
| Gumboro Vaccine     | 14 <sup>th</sup> day | 1 drop in each eye                                  |
| BCRDV               | 21 <sup>st</sup> day | 1 drop in each eye                                  |

### 3.6.7 Medication

An antibiotic named Docolis (0.5g/liter of drinking water) used for first 3 days to prevent early chick mortality. Eskavit WS (0.2g/liter drinking water), Electrocure + (0.5g/liter of drinking water) were given to prevent strives.

### 3.6.8 Sanitation

Proper hygienic measures and sanitation program of the experimental houses was taken during experimental period.

### 3.7 Post-mortem examination

Dead birds were diagnosed promptly at the Dairy and Poultry Science Laboratory of Hajee Mohammad Danesh Science and Technology University, Dinajpur. After postmortem examination, the result were collected and necessary measures were taken to solve the problem without applying medicines.



Fig 4: Postmortem examination of birds during experimental period.



### 3.8 Data collection and record keeping.

The following data were kept during the 5 weeks of rearing period:


- a. Body weight: Initial and at the end of each week.
- b. Feed consumption: Weekly
- c. Feed conversion ratio: weekly
- d. Survivability: Recorded from mortality.
- e. Dressing yield: At the end of experiment two broiler were slaughtered from each replication to estimate dressing yield including dressed weight, thigh weight, Drum stick weight, breast meat weight, wing meat weight, head weight, gizzard weight, heart weight, spleen weight, shank weight, abdominal fat weight, skin weight, viscera weight.
- f. Cost Benefit ratio.



Fig 5: Processing of birds during experimental period.

### 3.9 Statistical analysis

All recorded and calculated data were statistically analyzed by using analysis of variance (ANOVA) technique by a computer using “MSTAT statistical software” in accordance with the principles of Randomized Complete Block Design (RCBD). Duncan’s Multiple Range Test was done to know the differences among the treatment means at 5% and 1% level of significance (Duncan, 1955)



**CHAPTER 4**  
**RESULTS AND DISCUSSION**

## CHAPTER 4

### RESULTS AND DISCUSSION

#### 4.1 Effect of Phytase enzyme on body weight gain of broilers

Tables 4.1 and 4.2 showed the growth performance as affected by level phytase enzyme supplementation during the different weekly intervals and the entire experimental period (2<sup>nd</sup> to 5<sup>th</sup> wks of age). The initial (1st week) body weight in all treatment groups were similar. Significant differences ( $P>0.05$ ) were found on body weight among different treatment groups at 5<sup>th</sup> weeks of age. The body weight of T<sub>1</sub>F (Fast Feather fed diet with 1gm phytase enzyme per kg Feed) was highest at 2<sup>nd</sup> weeks of age. The body weight of T<sub>0</sub>F (Fast Feather fed diet without phytase enzyme) was more than T<sub>1</sub>A, T<sub>1</sub>H, T<sub>1</sub>C at 2<sup>nd</sup> & 3<sup>rd</sup> weeks of age. The highest body weight and body weight gain was attained by the birds of T<sub>1</sub>F (Fast feather fed diet with 1gm phytase enzyme per kg Feed) and T<sub>1</sub>C (Cobb-500 fed diet with 1 gm phytase enzyme per kg Feed) at 5<sup>th</sup> weeks of age. Among the treatment groups significant differences ( $P<0.01$ ) were found at T<sub>1</sub>H in comparison with T<sub>1</sub>C, T<sub>1</sub>F and T<sub>1</sub>A. The body weight of T<sub>1</sub>C, T<sub>1</sub>F and T<sub>1</sub>A (5<sup>th</sup> week) were similar. All the strain of treatment groups showed better body weight than control groups. Among the phytase group the result was significantly differed ( $P>0.05$ ). The highest body weight was found in T<sub>1</sub>C (1705.07g) and lowest in T<sub>1</sub>H (1589.55). In comparison between treatment and control groups, all strain of treatment groups showed significantly higher body weight than the control groups. The body weight of different treatments groups were T<sub>1</sub>C (1705.076g), T<sub>1</sub>F (1702.49g), T<sub>1</sub>A (1681.54g), T<sub>1</sub>H (1589.55g) and control groups were T<sub>0</sub>F (1650.0g), T<sub>0</sub>C (1624.25g), T<sub>0</sub>A (1526.10g), T<sub>0</sub>H (1468.06g) respectively. The result was consistent with the findings of some earlier workers (Qian *et al.*, 1996; Sebastian *et al.*, 1996; Nadeem *et al.*, 2005). They reported that body weight and body weight gain were increased in chicks due to fed phytase diet and utilization of p-from the phytase mineral complex. These results are in disagreement with a number of authors (Ravindran *et al.*, 2001; Kwon *et al.*, 1999; Sohail and Roland, 1999). Similar result was also obtained by Sebastian *et al.* (1997), however, phytase supplementation had no influence on the apparent ileal digestibility in their study.

**Table : 4.1** Effect of Phytase enzyme in the diet of broilers on body weight (g/bird )

| Age (Week)      | Dietary treatment groups      |                                |                                |                              |                                |                               |                                |                               |    |  |  | Level of Significance |
|-----------------|-------------------------------|--------------------------------|--------------------------------|------------------------------|--------------------------------|-------------------------------|--------------------------------|-------------------------------|----|--|--|-----------------------|
|                 | T <sub>0</sub> F              | T <sub>0</sub> A               | T <sub>0</sub> C               | T <sub>0</sub> H             | T <sub>1</sub> F               | T <sub>1</sub> A              | T <sub>1</sub> C               | T <sub>1</sub> H              |    |  |  |                       |
| Initial weight  | 41.67<br>0.13                 | 40.17<br>0.13                  | 39.83<br>0.13                  | 37.50<br>0.23                | 41.67<br>0.13                  | 40.83<br>0.49                 | 40.0<br>0.47                   | 39.67<br>0.27                 | NS |  |  |                       |
| 1 <sup>st</sup> | 151.8<br>3.018                | 132.3<br>0.906                 | 137.5<br>1.752                 | 125.4<br>2.264               | 151.7<br>2.744                 | 133.2<br>1.549                | 139.9<br>1.199                 | 127.5<br>1.732                | NS |  |  |                       |
| 2 <sup>nd</sup> | 339.2<br>6.182 <sup>ab</sup>  | 300.9<br>5.216 <sup>d</sup>    | 327.5<br>2.364 <sup>abc</sup>  | 293.9<br>2.551 <sup>d</sup>  | 353.6<br>5.308 <sup>a</sup>    | 313.4<br>4.768 <sup>bed</sup> | 334.2<br>1.732 <sup>abc</sup>  | 309.1<br>3.149 <sup>cd</sup>  | ** |  |  |                       |
| 3 <sup>rd</sup> | 690.2<br>1.433 <sup>b</sup>   | 586.7<br>3.485 <sup>d</sup>    | 605.2<br>3.054 <sup>d</sup>    | 604.8<br>4.267 <sup>d</sup>  | 746.1<br>10.665 <sup>a</sup>   | 637.7<br>5.197 <sup>c</sup>   | 633.5<br>5.667 <sup>c</sup>    | 637.1<br>3.018 <sup>c</sup>   | ** |  |  |                       |
| 4 <sup>th</sup> | 1196.35<br>8.665 <sup>b</sup> | 1065.43<br>5.247 <sup>cd</sup> | 1173.04<br>1.874 <sup>b</sup>  | 1017.17<br>3.00 <sup>d</sup> | 1317.38<br>14.423 <sup>a</sup> | 1176.41<br>5.215 <sup>b</sup> | 1230.00<br>22.671 <sup>b</sup> | 1087.00<br>3.389 <sup>c</sup> | ** |  |  |                       |
| 5 <sup>th</sup> | 1650.0<br>0.13 <sup>bc</sup>  | 1526.10<br>9.28 <sup>e</sup>   | 1624.25<br>10.36 <sup>cd</sup> | 1468.06<br>1.86 <sup>f</sup> | 1702.49<br>8.32 <sup>a</sup>   | 1681.54<br>3.95 <sup>a</sup>  | 1705.076<br>8.70 <sup>a</sup>  | 1589.55<br>2.05 <sup>d</sup>  | ** |  |  |                       |

Values of different variables under different program indicates Mean SEM ; abcdefg mean values with dissimilar super scripts are significantly different; SEM , Standards errors of means ; \*\* Significant (P<0.01) ; \* Significant (P>0.05); NS (Non-significant)

**Here,**

T<sub>0</sub>F : Fast feather + giving diet without phytase enzyme.

T<sub>0</sub>A : Arbor acres + giving diet without phytase enzyme.

T<sub>0</sub>C : Coob-500 + giving diet without phytase enzyme.

T<sub>0</sub>H : Hubbard classic + giving diet without phytase enzyme.

T<sub>1</sub>F : Fast feather + giving diet having 1g phytase enzyme/kg feed.

T<sub>1</sub>A : Arbor acres + giving diet having 1g phytase enzyme/kg feed.

T<sub>1</sub>C : Cobb-500 + giving diet having 1g phytase enzyme/kg feed.

T<sub>1</sub>H : Hubbard classic + giving diet having 1g phytase enzyme/kg feed.

**Table : 4.2** Effect of Phytase enzyme on the production performance of broiler.

| Parameters                     | Age in (Week)     | Dietary treatment groups |                    |                     |                    |                    |                    |                    |                    |    |  | Level Significance |
|--------------------------------|-------------------|--------------------------|--------------------|---------------------|--------------------|--------------------|--------------------|--------------------|--------------------|----|--|--------------------|
|                                |                   | T <sub>0</sub> F         | T <sub>0</sub> A   | T <sub>0</sub> C    | T <sub>0</sub> H   | T <sub>1</sub> F   | T <sub>1</sub> A   | T <sub>1</sub> C   | T <sub>1</sub> H   |    |  |                    |
| Body weight (g/bird)           | 0-5 <sup>th</sup> | 1650.0                   | 1526.10            | 1624.25             | 1468.06            | 1702.49            | 1681.54            | 1705.076           | 1589.55            | ** |  |                    |
|                                |                   | 0.13 <sup>bc</sup>       | 9.28 <sup>c</sup>  | 10.36 <sup>cd</sup> | 1.86 <sup>f</sup>  | 8.32 <sup>a</sup>  | 3.95 <sup>a</sup>  | 8.70 <sup>a</sup>  | 2.05 <sup>d</sup>  |    |  |                    |
| Body weight gain (g/bird)      | 0-5 <sup>th</sup> | 1608.31                  | 1485.93            | 1584.41             | 1428.06            | 1661.72            | 1640.70            | 1665.28            | 1549.88            | ** |  |                    |
|                                |                   | 12.84 <sup>abc</sup>     | 9.30 <sup>d</sup>  | 10.24 <sup>bc</sup> | 1.66 <sup>e</sup>  | 8.20 <sup>a</sup>  | 4.22 <sup>ab</sup> | 8.87 <sup>a</sup>  | 1.96 <sup>cd</sup> |    |  |                    |
| Feed consumption (g/bird)      | 0-5 <sup>th</sup> | 3029.205                 | 2788.042           | 2883.48             | 2656.12            | 2940.48            | 2826.78            | 2914.42            | 2638.47            | *  |  |                    |
|                                |                   | 12.84 <sup>abc</sup>     | 18.79 <sup>c</sup> | 16.70 <sup>bc</sup> | 20.20 <sup>d</sup> | 8.20 <sup>ab</sup> | 4.22 <sup>ab</sup> | 8.70 <sup>ab</sup> | 14.03 <sup>d</sup> |    |  |                    |
| Feed conversion ratio (g/bird) | 0-5 <sup>th</sup> | 1.88                     | 1.87               | 1.82                | 1.86               | 1.77               | 1.72               | 1.75               | 1.70               | ** |  |                    |
|                                |                   | 0.01 <sup>a</sup>        | 0.02 <sup>a</sup>  | 0.00 <sup>ab</sup>  | 0.01 <sup>a</sup>  | 0.01 <sup>bc</sup> | 0.01 <sup>c</sup>  | 0.01 <sup>bc</sup> | 7.20 <sup>c</sup>  |    |  |                    |
| Livability (%)                 | 0-5 <sup>th</sup> | 94.87                    | 94.87              | 97.43               | 92.30              | 97.43              | 97.43              | 97.43              | 97.43              | NS |  |                    |
|                                |                   | 4.18                     | 4.18               | 2.09                | 3.62               | 2.09               | 2.09               | 2.09               | 2.09               |    |  |                    |
| Profit / broiler (Tk.)         | 0-5 <sup>th</sup> | 25.09                    | 20.64              | 33                  | 14                 | 35.43              | 37.87              | 36                 | 34                 |    |  |                    |
|                                |                   |                          |                    | .88                 | .55                |                    |                    | .96                | .69                |    |  |                    |

Values of different variables under different program indicates Mean SEM ; abcdefg mean values with dissimilar super scripts a significantly different; SEM , Standards errors of means ; \*\* Significant (P<0.01) ; \* Significant (P>0.05); NS (Non-significant)

**Here,**

T<sub>0</sub>F : Fast feather + giving diet without phytase enzyme.

T<sub>0</sub>A : Arbor acres + giving diet without phytase enzyme.

T<sub>0</sub>C : Coob-500 + giving diet without phytase enzyme.

T<sub>0</sub>H : Hubbard classic + giving diet without phytase enzyme.

T<sub>1</sub>F : Fast feather + giving diet having 1g phytase enzyme/kg feed.

T<sub>1</sub>A : Arbor acres + giving diet having 1g phytase enzyme/kg feed.

T<sub>1</sub>C : Cobb-500 + giving diet having 1g phytase enzyme/kg feed.

T<sub>1</sub>H : Hubbard classic + giving diet having 1g phytase enzyme/kg feed.

## **4.2 Effect of Phytase enzyme on feed intake of broilers**

The differences in feed intake between control groups (birds fed diet without phytase enzyme/kg feed) and treatment groups (birds fed diet with 1 gm phytase enzyme) were significant ( $P>0.05$ ) at 5<sup>th</sup> weeks of age. The result on feed intake (Table 4.2 and 4.3) demonstrates that during 28-35 days of age the birds in T<sub>0</sub>F (Fast feather fed diet without phytase enzyme) consumed more feed than T<sub>1</sub>F, T<sub>0</sub>C, T<sub>1</sub>C, T<sub>0</sub>A, T<sub>1</sub>A, T<sub>0</sub>H and T<sub>1</sub>H consequently. Among the treatment groups significant differences ( $P<0.05$ ) were found at T<sub>1</sub>H in comparison with T<sub>1</sub>F, T<sub>1</sub>C and T<sub>1</sub>A. The feed consumption of T<sub>1</sub>F, T<sub>1</sub>C, T<sub>1</sub>A (5<sup>th</sup> week) were similar. All the strain of treatment groups consumed less food than the control groups. The feed consumption of different treatment groups were T<sub>1</sub>F (2940.48g), T<sub>1</sub>C (2914.42g), T<sub>1</sub>A (2826.78g), T<sub>1</sub>H (2638.47g) and control groups were T<sub>0</sub>F (3029.205g), T<sub>0</sub>C (2883.042g), T<sub>0</sub>A (2788.042g), T<sub>0</sub>H (2656.12g) respectively. Among the treatment groups the feed intake of broilers were significantly differed ( $P>0.05$ ). The highest feed intake was found in T<sub>1</sub>F (2940.48g) and lowest in T<sub>1</sub>H (2638.47). In comparison between treatment and control groups, all strain of treatment groups showed lower feed intake than control groups. The result was consistent with the findings of some earlier workers (Wilson *et al.*, 1999). They found that feed consumption was decreased due to addition of enzyme since birds fulfilled their nutrient requirement by taking less amount of feed. These result inconsistent with Ahmad *et al.*, (2000) who reported that in addition of phytase did not affect feed consumption.

## **4.3 Effect of Phytase enzyme on feed conversion ratio (FCR) of broilers**

The feed conversion ratio between control groups (Birds fed diet without phytase enzyme/kg feed) and treatment groups (Birds fed diet with 1 gm phytase enzyme) were significantly different ( $P<0.01$ ). Table 4.2 and 4.4 showed that significant differences ( $P>0.01$ ) were found on feed conversion ratio at 5<sup>th</sup> weeks of age. Results showed that among the four different strains, Hubbard classic had the lowest FCR, suggesting that this broiler strain could utilize the feed efficiently compared to other three strains. All the strain of treatment groups showed better FCR than control groups. The feed conversion ratio of different treatment groups were T<sub>1</sub>H (1.70), T<sub>1</sub>A

(1.72), T<sub>1</sub>C (1.75), T<sub>1</sub>F (1.77) and control groups were T<sub>0</sub>C (1.82), T<sub>0</sub>H (1.86), T<sub>0</sub>A (1.87), T<sub>0</sub>F (1.88) respectively. At the end of trail, the feed conversion ratio of T<sub>1</sub>H (Hubbard classic fed diet with 1gm phytase enzyme/kg feed) was better followed by T<sub>1</sub>A, T<sub>1</sub>C, T<sub>1</sub>F, T<sub>0</sub>C, T<sub>0</sub>H, T<sub>0</sub>A, T<sub>0</sub>F, respectively. The addition of exogenous phytase improved ( $P < 0.05$ ) feed conversion ratio (Table 4.4) of the birds. In comparison between treatment and control groups, all strain of treatment groups significantly ( $P > 0.05$ ) lower feed conversion ratio than control groups. The improvement of feed conversion ratio of chicks as a result of phytase application was in agreement with the results of some earlier studies (Shirley *et al.*, 2003; Selle *et al.*, 2006; Watson *et al.*, 2006). In those studies, feed conversion ratio of different broiler groups differed significantly might be due to the increased availability of energy and amino acid digestibility by raise in phytate degradation. The improve of body weight gain and feed conversion ratio of chicks as a result of phytase application might be due to the increased availability of energy (Shirley and Edwards, 2003) and amino acid digestibility (Selle *et al.*, 2006) due to a raise in phytate degradation. The ability of phytase to improve P availability by hydrolyzing phytate-bound P in poultry diets can therefore reduce supplementation of diets with inorganic P sources.

**Table : 4.3** Effect of Phytase enzyme in the diet of broilers on feed intake (g/bird)

| Age (Week)      | Dietary treatment groups |                      |                     |                     |                     |                      |                     |                      |                      |                      | Level of Significance |
|-----------------|--------------------------|----------------------|---------------------|---------------------|---------------------|----------------------|---------------------|----------------------|----------------------|----------------------|-----------------------|
|                 | T <sub>0</sub> F         | T <sub>0</sub> A     | T <sub>0</sub> C    | T <sub>0</sub> H    | T <sub>1</sub> F    | T <sub>1</sub> A     | T <sub>1</sub> C    | T <sub>1</sub> H     |                      |                      |                       |
| 1 <sup>st</sup> | 151.7                    | 129.2                | 140.3               | 122.8               | 150.4               | 129.0                | 141.9               | 122.5                | 122.5                | 3.389                | NS                    |
|                 | 5.286                    | 1.615                | 2.005               | 2.852               | 3.714               | 1.945                | 2.460               | 3.389                | 3.389                |                      |                       |
| 2 <sup>nd</sup> | 460.8                    | 399.0                | 437.2               | 397.7               | 445.1               | 394.2                | 442.4               | 388.7                | 388.7                | 1.971                | NS                    |
|                 | 5.006                    | 10.711               | 4.299               | 2.357               | 9.775               | 6.786                | 6.950               | 1.971                | 1.971                |                      |                       |
| 3 <sup>rd</sup> | 1076.40                  | 896.8                | 915.8               | 947.9               | 1066.11             | 887.3                | 919.7               | 911.6                | 911.6                | 12.854 <sup>dc</sup> | *                     |
|                 | 8.509 <sup>a</sup>       | 10.192 <sup>bc</sup> | 5.858 <sup>bc</sup> | 9.797 <sup>d</sup>  | 18.188 <sup>a</sup> | 5.765 <sup>c</sup>   | 3.598 <sup>dc</sup> | 12.854 <sup>dc</sup> | 12.854 <sup>dc</sup> |                      |                       |
| 4 <sup>th</sup> | 2059.66                  | 1808.42              | 1949.06             | 1743.44             | 2084.75             | 18.09.17             | 1918.42             | 1713.41              | 1713.41              | 3.281 <sup>d</sup>   | *                     |
|                 | 4.296 <sup>a</sup>       | 15.521 <sup>c</sup>  | 7.762 <sup>d</sup>  | 6.021 <sup>cd</sup> | 32.331 <sup>a</sup> | 11.1116 <sup>c</sup> | 27.598 <sup>b</sup> | 3.281 <sup>d</sup>   | 3.281 <sup>d</sup>   |                      |                       |
| 5 <sup>th</sup> | 3029.205                 | 2788.042             | 2883.48             | 2656.12             | 2940.48             | 2826.78              | 2914.42             | 2638.47              | 2638.47              | 14.03 <sup>d</sup>   | *                     |
|                 | 33.88 <sup>a</sup>       | 18.79 <sup>c</sup>   | 16.70 <sup>bc</sup> | 20.20 <sup>d</sup>  | 8.20 <sup>ab</sup>  | 4.22 <sup>ab</sup>   | 8.70 <sup>ab</sup>  | 14.03 <sup>d</sup>   | 14.03 <sup>d</sup>   |                      |                       |

Values of different variables under different program indicates Mean SEM ; abcdefg mean values with dissimilar super scripts are significantly different; SEM , Standards errors of means ; \*\* Significant (P<0.01) ; \* Significant (P>0.05); NS (Non-significant)

**Here,**

T<sub>0</sub>F : Fast feather + giving diet without phytase enzyme.

T<sub>0</sub>A : Arbor acres + giving diet without phytase enzyme.

T<sub>0</sub>C : Coob-500 + giving diet without phytase enzyme.

T<sub>0</sub>H : Hubbard classic + giving diet without phytase enzyme.

T<sub>1</sub>F : Fast feather + giving diet having 1g phytase enzyme/kg feed.

T<sub>1</sub>A : Arbor acres + giving diet having 1g phytase enzyme/kg feed.

T<sub>1</sub>C : Cobb-500 + giving diet having 1g phytase enzyme/kg feed.

T<sub>1</sub>H : Hubbard classic + giving diet having 1g phytase enzyme/kg feed.



**Table : 4.4** Effect of Phytase enzyme in the diet of broilers on feed conversion ratio (g/bird)

| Age (Week)      | Dietary treatment groups   |                             |                             |                            |                            |                             |                              |                             |  |    | Level of Significance |
|-----------------|----------------------------|-----------------------------|-----------------------------|----------------------------|----------------------------|-----------------------------|------------------------------|-----------------------------|--|----|-----------------------|
|                 | T <sub>0</sub> F           | T <sub>0</sub> A            | T <sub>0</sub> C            | T <sub>0</sub> H           | T <sub>1</sub> F           | T <sub>1</sub> A            | T <sub>1</sub> C             | T <sub>1</sub> H            |  |    |                       |
| 1 <sup>st</sup> | 1.37<br>0.011              | 1.40<br>0.005               | 1.43<br>0.016               | 1.43<br>0.009              | 1.36<br>0.007              | 1.39<br>0.007               | 1.42<br>0.12                 | 1.39<br>0.011               |  | NS |                       |
| 2 <sup>nd</sup> | 1.54<br>0.016 <sup>a</sup> | 1.52<br>0.020 <sup>ab</sup> | 1.51<br>0.016 <sup>ab</sup> | 1.55<br>0.005 <sup>a</sup> | 1.42<br>0.004 <sup>c</sup> | 1.44<br>0.012 <sup>bc</sup> | 1.48<br>0.012 <sup>abc</sup> | 1.43<br>0.022 <sup>bc</sup> |  | ** |                       |
| 3 <sup>rd</sup> | 1.65<br>0.016 <sup>a</sup> | 1.63<br>0.015 <sup>a</sup>  | 1.61<br>0.014 <sup>ab</sup> | 1.67<br>0.009 <sup>a</sup> | 1.50<br>0.005 <sup>c</sup> | 1.47<br>0.007 <sup>c</sup>  | 1.54<br>0.009 <sup>bc</sup>  | 1.51<br>0.017 <sup>c</sup>  |  | ** |                       |
| 4 <sup>th</sup> | 1.77<br>0.016 <sup>a</sup> | 1.75<br>0.023 <sup>a</sup>  | 1.71<br>0.009 <sup>a</sup>  | 1.77<br>0.009 <sup>a</sup> | 1.62<br>0.011 <sup>b</sup> | 1.58<br>0.014 <sup>b</sup>  | 1.60<br>0.007 <sup>b</sup>   | 1.62<br>0.007 <sup>b</sup>  |  | ** |                       |
| 5 <sup>th</sup> | 1.88<br>0.01 <sup>a</sup>  | 1.87<br>0.02 <sup>a</sup>   | 1.82<br>0.00 <sup>a</sup>   | 1.86<br>0.01 <sup>a</sup>  | 1.77<br>0.01 <sup>b</sup>  | 1.72<br>0.01 <sup>b</sup>   | 1.75<br>0.01 <sup>b</sup>    | 1.70<br>0.007 <sup>c</sup>  |  | ** |                       |

Values of different variables under different program indicates Mean SEM ; abcdefg mean values with dissimilar super scripts are significantly different; SEM , Standards errors of means ; \*\* Significant (P<0.01) ; \* Significant (P>0.05); NS (Non-significant)

**Here,**

T<sub>0</sub>F : Fast feather + giving diet without phytase enzyme.

T<sub>0</sub>A : Arbor acres + giving diet without phytase enzyme.

T<sub>0</sub>C : Coob-500 + giving diet without phytase enzyme.

T<sub>0</sub>H : Hubbard classic + giving diet without phytase enzyme.

T<sub>1</sub>F : Fast feather + giving diet having 1g phytase enzyme/kg feed.

T<sub>1</sub>A : Arbor acres + giving diet having 1g phytase enzyme/kg feed.

T<sub>1</sub>C : Cobb-500 + giving diet having 1g phytase enzyme/kg feed.

T<sub>1</sub>H : Hubbard classic + giving diet having 1g phytase enzyme/kg feed.

#### **4.4 Effect of Phytase enzyme on carcass characteristics of broilers**

The meat yield characteristics are shown in Table 4.5. The result indicates that there was no significant differences among the treatments for breast meat weight, wing meat weight, head weight, gizzard weight, heart weight, spleen weight, shank weight, abdominal fat weight, skin weight and viscera weight. On the other hand, the dressing yield, thigh weight and drum stick weight was significantly different ( $P < 0.01$ ) between treatment groups (Birds fed diet with 1gm phytase enzyme/kg feed) and control groups (Birds fed diet without phytase enzyme) during 5 weeks of experimental period. The dressing yield, drumstick weight and thigh weight were higher in T<sub>1</sub>H, T<sub>1</sub>C and T<sub>1</sub>F. The increased dressing yield, drum stick weight and thigh weight on T<sub>1</sub>H, T<sub>1</sub>C, T<sub>1</sub>F might be because of the increased live weight. But these findings contradict with the observation of Bharathidhasan *et al.* (2009) who found that a marginal increase in dressing percentage in birds fed with diets containing enzyme level at 0, 250, 500, 750 and 1,000 g ton<sup>-1</sup> of feed. Also, Jordão Filho *et al.* (2006) did not find differences using 500 until 1,500 FTU phytase kg<sup>-1</sup> of feed. These results also agreed with previous findings of Angel *et al.*, (2007) but opposite to those of Pillai *et al.*, (2006) who showed that phytase supplementation significantly increased percentages of most of carcass merits compared to P-deficient diets.

#### **4.5 Effect of Phytase enzyme on production cost and profit margin of broilers**

Phytase supplementation, by improving overall production performance of broilers, may lead to economic benefits. The production cost of broiler was shown in Table 4.6. The feed cost was highest (Tk. 5316) in treatment T<sub>0</sub>F (Control group) and the lowest (Tk. 4630) in treatment T<sub>1</sub>H (Treatment group). The opposite result was found in T<sub>1</sub>C and T<sub>1</sub>A (Treatment groups) where the addition of phytase enzyme resulted in increased feed cost in treatment T<sub>0</sub>C and T<sub>0</sub>A (Control groups). The cost per kg live weight of broiler was the highest in treatment T<sub>1</sub>F (Tk. 180.15) followed by treatment T<sub>1</sub>C (179.0), T<sub>0</sub>F (178.41), T<sub>1</sub>A (175.05), T<sub>0</sub>C (171.82), T<sub>0</sub>A (167.56), T<sub>1</sub>H (166.58) & T<sub>0</sub>H (161.61) respectively. Net profit per live broiler was the highest in treatment T<sub>1</sub>A (Tk. 37.87) followed by treatment T<sub>1</sub>C (36.96), T<sub>1</sub>F (35.43), T<sub>1</sub>H (34.69), T<sub>0</sub>C

(33.88), T<sub>0</sub>F (25.09), T<sub>0</sub>A (20.64) and T<sub>0</sub>H (14.55) respectively. The highest net profit was observed in treatment T<sub>1</sub>A (Arber acres fed on diet treated with 1gm phytase enzyme/kg feed) and the lowest total net profit was observed in treatment T<sub>0</sub>H (Habbard classic fed on diet without phytase enzyme). Economic wise, the addition of phytase could make reasonable profits than without its addition. The findings of the present study is similar to the observation of Islam *et al.* (2010). They reported that net profits (Kg live bird) was significantly better in broiler group fed diet supplemented with enzyme 50g/100kg which might be due to improvement of digestibility's and consequent by better utilization of nutrients. The profitability element here is based on feed, as it constitutes more than 60% of the ration cost for poultry feeding. The result was consistent with the result of Augelovicova and Michalik (1997) who stated that enzyme supplementation in commercial broiler diet decreased feed cost by 8.81% to 9.73% for production of 1 kg broiler meat. Vinil *et al.* (2000) found a reduction in feed cost in soy-wheat bran diets supplemented with phytase (25 g 100 kg<sup>-1</sup>) of about 1.00 Indian rupee (INR). Net income increased up to 9.47% in response to 300 g/ton of phytase supplementa-tion (Kundu *et al.* 2000). Singh and Khatta (2004) reported that phytase supplementation resulted in 10% and 6% re-ductions in cost per unit gain in broilers fed corn and wheat based diets, respectively. Plumstead *et al.* (2008) observed that less expensive broiler diets low in P and other nutrients supplemented with phytase resulted in optimum production. Supplementation of phytase leads to safe, economic and almost complete replacement of dietary P (dicalcium phos-phate), that ultimately causes reduction in feed cost kg<sup>-1</sup> of weight gain (Singh *et al.* 2003a, b; Singh and Khatta, 2003a, b).

**Table : 4.5** Effect of Phytase enzyme in the diet of broilers on carcass characteristics at 35 days of age.

| Parameters           | Dietary treatment groups     |                              |                                |                              |                              |                               |                               |                              |  |  | Level of Significance |
|----------------------|------------------------------|------------------------------|--------------------------------|------------------------------|------------------------------|-------------------------------|-------------------------------|------------------------------|--|--|-----------------------|
|                      | T <sub>0</sub> F             | T <sub>0</sub> A             | T <sub>0</sub> C               | T <sub>0</sub> H             | T <sub>1</sub> F             | T <sub>1</sub> A              | T <sub>1</sub> C              | T <sub>1</sub> H             |  |  |                       |
| Live wt (g/bird)     | 1650.0<br>0.13 <sup>bc</sup> | 1526.10<br>9.28 <sup>e</sup> | 1624.25<br>10.36 <sup>cd</sup> | 1468.06<br>1.86 <sup>f</sup> | 1702.49<br>8.32 <sup>a</sup> | 1681.54<br>3.95 <sup>ab</sup> | 1705.076<br>8.70 <sup>a</sup> | 1589.55<br>2.05 <sup>d</sup> |  |  | **                    |
| Dressing wt (%)      | 81.32<br>0.72 <sup>e</sup>   | 84.61<br>0.22 <sup>d</sup>   | 87.75<br>0.33 <sup>bc</sup>    | 85.71<br>0.47 <sup>cd</sup>  | 91.24<br>0.17 <sup>a</sup>   | 90.38<br>0.98 <sup>ab</sup>   | 91.30<br>0.41 <sup>a</sup>    | 92.19<br>0.35 <sup>a</sup>   |  |  | **                    |
| Drum stick wt (%)    | 8.390<br>0.188 <sup>d</sup>  | 8.073<br>0.242 <sup>b</sup>  | 8.407<br>0.119 <sup>b</sup>    | 8.883<br>0.035 <sup>ab</sup> | 9.450<br>0.258 <sup>a</sup>  | 9.913<br>0.152 <sup>a</sup>   | 9.783<br>0.034 <sup>a</sup>   | 8.953<br>0.205 <sup>ab</sup> |  |  | **                    |
| Thigh wt (%)         | 11.60<br>0.28 <sup>b</sup>   | 11.36<br>0.17 <sup>bc</sup>  | 10.60<br>0.26 <sup>c</sup>     | 10.86<br>0.19 <sup>bc</sup>  | 12.77<br>0.072 <sup>a</sup>  | 11.63<br>0.115 <sup>b</sup>   | 12.57<br>0.024 <sup>a</sup>   | 11.09<br>0.136 <sup>bc</sup> |  |  | **                    |
| Breast meat wt (%)   | 18.61<br>0.284               | 18.61<br>0.076               | 17.83<br>0.198                 | 18.72<br>0.145               | 18.29<br>0.143               | 18.40<br>0.229                | 18.60<br>0.086                | 19.07<br>0.153               |  |  | NS                    |
| Wing meat wt (%)     | 8.300<br>0.11                | 8.280<br>0.339               | 8.590<br>0.132                 | 8.287<br>0.056               | 8.620<br>0.263               | 8.570<br>0.043                | 8.400<br>0.143                | 8.530<br>0.251               |  |  | NS                    |
| Head wt (%)          | 2.337<br>0.066               | 2.363<br>0.040               | 2.187<br>0.051                 | 2.493<br>0.050               | 2.110<br>0.036               | 2.067<br>0.025                | 2.047<br>0.035                | 2.140<br>0.047               |  |  | NS                    |
| Gizzard wt (%)       | 1.883<br>0.107               | 1.967<br>0.080               | 2.037<br>0.089                 | 2.153<br>0.040               | 1.633<br>0.056               | 1.733<br>0.058                | 1.743<br>0.055                | 1.940<br>0.033               |  |  | NS                    |
| Heart wt (%)         | 0.55<br>0.012                | 0.56<br>0.008                | 0.536<br>0.016                 | 0.57<br>0.008                | 0.50<br>0.009                | 0.52<br>0.019                 | 0.52<br>0.007                 | 0.496<br>0.002               |  |  | NS                    |
| Spleen wt (%)        | 0.13<br>0.005                | 0.14<br>0.007                | 0.13<br>0.00                   | 0.15<br>0.002                | 0.13<br>0.004                | 0.11<br>0.002                 | 0.13<br>0.007                 | 0.13<br>0.002                |  |  | NS                    |
| Shank wt (%)         | 4.33<br>0.117                | 4.23<br>0.079                | 4.13<br>0.10                   | 4.31<br>0.119                | 4.95<br>0.172                | 4.63<br>0.073                 | 4.80<br>0.069                 | 4.14<br>0.026                |  |  | NS                    |
| Abdominal Fat wt (%) | 1.79<br>0.086                | 1.76<br>0.032                | 1.66<br>0.087                  | 2.08<br>0.039                | 1.46<br>0.032                | 1.34<br>0.024                 | 1.45<br>0.034                 | 1.67<br>0.036                |  |  | NS                    |
| Skin wt (%)          | 6.51<br>0.112                | 6.34<br>0.061                | 6.00<br>0.108                  | 7.16<br>0.116                | 6.86<br>0.115                | 6.45<br>0.033                 | 6.73<br>0.036                 | 6.71<br>0.168                |  |  | NS                    |
| Viscera wt (%)       | 10.20<br>0.208               | 10.23<br>0.198               | 9.79<br>0.026                  | 11.27<br>0.2506              | 9.25<br>0.103                | 9.26<br>0.088                 | 9.55<br>0.150                 | 10.16<br>0.019               |  |  | NS                    |

Here,

00T<sub>0</sub>F : Fast feather + giving diet without phytase enzyme.

T<sub>0</sub>A : Arbor acres + giving diet without phytase enzyme.

T<sub>0</sub>C : Coob-500 + giving diet without phytase enzyme.

T<sub>0</sub>H : Hubbard classic + giving diet without phytase enzyme.

T<sub>1</sub>F : Fast feather + giving diet having 1g phytase enzyme/kg feed.

T<sub>1</sub>A : Arbor acres + giving diet having 1g phytase enzyme/kg feed.

T<sub>1</sub>C : Coob-500 + giving diet having 1g phytase enzyme/kg feed.

T<sub>1</sub>H : Hubbard classic + giving diet having 1g phytase enzyme/kg feed.

**Table : 4.6** Effect of Phytase enzyme on production cost and profit margin of broilers.

| Cost of production (TK)    | Dietary treatment groups |                  |                  |                  |                  |                  |                  |                  |  |  |  |
|----------------------------|--------------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|--|--|--|
|                            | T <sub>0</sub> F         | T <sub>0</sub> A | T <sub>0</sub> C | T <sub>0</sub> H | T <sub>1</sub> F | T <sub>1</sub> A | T <sub>1</sub> C | T <sub>1</sub> H |  |  |  |
| Feed cost @45tk/kg         | 5316                     | 4893             | 5 059            | 4 661            | 5159             | 4 960            | 5 114            | 4 630            |  |  |  |
| Chick cost @32tk/bird      | 1248                     | 1248             | 1 248            | 1 248            | 1248             | 1 248            | 1 248            | 1 248            |  |  |  |
| Disinfectant cost          | 10                       | 10               | 10 10            |                  | 10               | 10 10            | 10               |                  |  |  |  |
| Vaccination cost           | 50                       | 50               | 50 50            |                  | 50               | 50 50            | 50               |                  |  |  |  |
| Medicine cost              | 24                       | 24               | 24 24            |                  | 24               | 24 24            | 24               |                  |  |  |  |
| Transport cost             | 10                       | 10               | 10 10            |                  | 10               | 10 10            | 10               |                  |  |  |  |
| Depreciation cost          | 150                      | 150              | 150 150          | 15 0             | 150              | 15 0             | 150              | 15 0             |  |  |  |
| Litter cost                | 50                       | 50               | 50 50            |                  | 50               | 50 50            | 50               |                  |  |  |  |
| Labour cost                | 60                       | 60               | 60 60            |                  | 60               | 60 60            | 60               |                  |  |  |  |
| Water and electricity cost | 40                       | 40               | 40 40            |                  | 40               | 40 40            | 40               |                  |  |  |  |
| Phytase enzyme cost        |                          |                  |                  |                  | 225              | 225              | 225              | 225              |  |  |  |
| <b>Total Cost</b>          | <b>6958</b>              | <b>6535</b>      | <b>6 701</b>     | <b>6 303</b>     | <b>7026</b>      | <b>6 827</b>     | <b>6 981</b>     | <b>6 497</b>     |  |  |  |
| Cost/broiler               | 178.41                   | 167.56           | 171.82           | 16 1.61          | 180.15           | 17 5.05          | 179              | 166.58           |  |  |  |
| Total selling @ 130tk/kg   | 7936.5                   | 7340.06          | 8022.56          | 6870.24          | 8407.88          | 8304.14          | 8422.7           | 7849.66          |  |  |  |
| Profit/broiler             | 25.09                    | 20.64            | 33 .88           | 14 .55           | 35.43            | 37 .87           | 36 .96           | 34 .69           |  |  |  |

**Here,**

T<sub>0</sub>F : Fast feather + giving diet without phytase enzyme.

T<sub>0</sub>A : Arbor acres + giving diet without phytase enzyme.

T<sub>0</sub>C : Coob-500 + giving diet without phytase enzyme.

T<sub>0</sub>H : Hubbard classic + giving diet without phytase enzyme.

T<sub>1</sub>F : Fast feather + giving diet having 1g phytase enzyme/kg feed.


T<sub>1</sub>A : Arbor acres + giving diet having 1g phytase enzyme/kg feed.

T<sub>1</sub>C : Cobb-500 + giving diet having 1g phytase enzyme/kg feed.

T<sub>1</sub>H : Hubbard classic + giving diet having 1g phytase enzyme/kg feed.

#### **4.6 Effect of Phytase enzyme on livability of broilers**

The livability during the experimental period was 97.43, 97.43, 97.43, 97.43, 97.43, 94.87, 94.87, and 92.30 percent for T<sub>1</sub>A, T<sub>0</sub>C, T<sub>1</sub>H, T<sub>1</sub>F, T<sub>1</sub>C, T<sub>0</sub>F, T<sub>0</sub>A and T<sub>0</sub>H respectively. No significant differences were observed in livability among different treatment groups and control groups during 5 weeks of experiment. Livability of broilers fed on phytase enzyme was very much acceptable during the study period. The result of present study was consistent with the findings of some earlier studies (Pillai *et al.*, 1995; Alam, 2001).



**CHAPTER 5**  
**SUMMARY AND CONCLUSIONS**

## CHAPTER 5

### SUMMARY AND CONCLUSION

The objectives of this experiment were to know the efficacy of supplementation of exogenous phytase enzyme on productivity and carcass characteristics of different strain of commercial broilers. There were 4 strains of commercial broiler such as Arbor acres, Fast feather, Cobb-500 and Hubbard classic each having 78 birds were used in the experiment for a period of 5 weeks. A total number of 312 day old straight run broiler chicks were distributed to two dietary treatments i.e. basal diet (Control-T<sub>0</sub>) and basal diet supplemented with phytase enzyme @ 1gm/kg feed. The birds were fed on ad libitum basis and fresh drinking water was available for all the times. Body weight, feed intake, feed conversion ratio, livability, production cost and profit margin and carcass yields of different strains of broilers of different treatments were recorded.

At 5<sup>th</sup> week of age, the final body weight of broilers at T<sub>0</sub>F, T<sub>0</sub>A, T<sub>0</sub>C, T<sub>0</sub>H, T<sub>1</sub>F, T<sub>1</sub>A, T<sub>1</sub>C and T<sub>1</sub>H treatment groups were 1650.00g, 1526.10g, 1624.25g, 1468.06g, 1702.49g, 1681.54g, 1705.076g and 1589.55g respectively. Live weight was increased significantly ( $P < 0.01$ ) on T<sub>1</sub>C (Cobb-500 fed diet with 1g phytase enzyme/kg feed) and T<sub>1</sub>F (Fast feather fed diet with 1g phytase enzyme/kg feed) compared to control. There were significant difference ( $P > 0.05$ ) among different treatments in relation to feed consumption. However, total feed consumption on T<sub>0</sub>F, T<sub>0</sub>A, T<sub>0</sub>C, T<sub>0</sub>H, T<sub>1</sub>F, T<sub>1</sub>A, T<sub>1</sub>C and T<sub>1</sub>H treatment groups were 3029.20g, 2788.04g, 2883.48g, 2656.12g, 2940.48g, 2826.78g, 2914.42g and 2638.47g respectively at 5<sup>th</sup> week of age. Significant differences ( $P < 0.01$ ) were found in feed conversion among birds fed on diet treated with phytase enzyme. Feed conversion ratio during the 5<sup>th</sup> week of age was 1.88, 1.87, 1.82, 1.86, 1.77, 1.72, 1.75 and 1.70 and T<sub>0</sub>F, T<sub>0</sub>A, T<sub>0</sub>C, T<sub>0</sub>H, T<sub>1</sub>F, T<sub>1</sub>A, T<sub>1</sub>C and T<sub>1</sub>H treatment groups respectively. No significant differences were observed in livability among the treatment groups. The cost of production was the highest in treatment T<sub>1</sub>F followed by treatment T<sub>1</sub>C, T<sub>0</sub>F, T<sub>1</sub>A, T<sub>0</sub>C, T<sub>0</sub>A, T<sub>1</sub>H, T<sub>0</sub>H. Net profit per live broiler was the highest in



treatment T<sub>1</sub>A followed by treatment T<sub>1</sub>C, T<sub>1</sub>F, T<sub>1</sub>H, T<sub>0</sub>C, T<sub>0</sub>F, T<sub>0</sub>A, T<sub>0</sub>H respectively. Dressing parameters were almost similar in different treatments and the differences were insignificant among treatment but the dressing weight percentage, thigh weight percentage and drumstick weight percentage were significant ( $P>0.01$ ). From this study, it can be concluded that addition of phytase enzyme had a positive effect on bird's performance i.e. supplementation of phytase enzyme can enhance the growth performance of broilers in term of body weight, weight gain, feed conversion efficiency at reduced cost of broilers. These effects appeared to be more pronounced in the thermos table phytase supplemented group. Costs of broilers rations might be reduced as commercial sources of P in rations are reduced and promoting meat quality of broiler chicken with considerable reduction in overall cost of production. However, more research is needed to support these findings.



# REFERENCES

## REFERENCES

- Ahmad, T., Rasool, S., Sarwar, M., Haq, A. and Hasan, Z. (2000). Effect of microbial phytase produced from a fungus *aspergillus niger* on bioavailability of phosphorus and calcium in broiler chickens. *Animal Feed Science and Technology*, 83(2): 103-114.
- Aksakal, N. and Bilal, H. (2000). Feed enzymes. *Animal Feed Science and Technology*, 45:65–79.
- Aksakal, N. and Bilal, H. (2002). Growth, Livability and feed conversion of 1957 versus 2001 broilers when fed representative 1957 and 2001 broiler diets. *Poultry Science*. 82:1500-1508.
- Alam, S.H. (2001). Effect of exogenous enzymes in diet on broiler performance. *International Journal of Poultry Science*, 2: 168-173.
- Angel, R., Saylor, W. W., Mitchell, A. D., Powers, W. and Applegate, T. J. (2007). Effect of dietary Phosphorous, phytase and 25-hydroxycholecalciferol on broiler chicken bone mineralization, nutrient utilization and excreta quality of broiler chickens, *Poultry Science*, 87:1200-1211.
- Aoyagi, S. and Baker, D.H. (1995). Effect of microbial phytase and 1, 25-dihydroxy xholecalciferol on dietary copper utilization in chicks. *Poultry Science*. 74:121-126.
- Atia, F.A., Waibel, P. E., Hermes, I., Carlson, C.W. and Walser, M. M. (2000). Effect of dietary phosphorus, calcium, and phytase on performance of growing turkeys. *Poultry Science*. 79:231–239.
- Augelovicova, M. and Michalik, I. (1997). A test of enzymatic preparation in relation to performance and commercial utilization of feeds in broiler chickens. *Zivocisna-Vyroba*, 42: 175-180.
- Aureli, R., Faruk, M. U., Cechova, I., Pedersen, P., Elvig-Joergensen, S., Fru, F. and Broz, J. (2011). The efficacy of a novel microbial 6-phytase expressed in *aspergillusoryzae* on the performance and phosphorus utilization in broiler chickens. *International Journal of Poultry Science*. 10: 160-168.

- Bharathidhasan, M. R., Luis, E. S. and Centeno, J. R. 2009. Performance of broilers fed corn-soybean based diets with different levels of enzyme and probiotic supplement. *Philippine Journal of Veterinary Medicine*, 39: 58-66.
- Biehl, R. R. and Baker, D. H. (1997). Microbial phytase improves amino acid utilization in young chicks fed diets based on soybean meal but not diets based on peanut meal. *Poultry Science*. 76:355-360.
- Broz, J., Oldale, P., Perrin-Voltz, A.H., Rychen, G., Schulze, J. and Simoes, C. (1994). Effect of supplemental phytase on performance and phosphorus utilization in broiler chickens fed a low phosphorus diet without addition of inorganic phosphates. *British Poultry Science*. 35(2): 273-280.
- Brenes, A., Viveros, A., Arija, I., Centeno, C., Pizarro, M. and Bravo, C. (2003). The effect of citric acid and microbial phytase on mineral utilization in broiler chicks. *Animal Feed Science and Technology*. 110: 201- 219.
- Caldwell, R.A. (1992). Effect of calcium and phytic acid on the activation of trypsinogen and the stability of trypsin. *Journal for Agriculture Food Chemistry*. 40: 43-46.
- Cheryan, M. and Rackis, J. J. (1980). Phytic acid interactions in food systems. *Crit. Rev. Food Sci*. 13: 297-335.
- Ciftci, M., Bestami, D. and Azman, M. (2005). Effects of microbial phytase supplementation on feed consumption and egg production of laying hens. *International Journal of Poultry Science*. 4 (10): 758- 760.
- Deshpande, S. S. and Cheryan, M. (1984). Effect of phytic acid, divalent cations and their interaction on alpha amylase activity. *Journal of Food Science*. 49:10
- Driver, J. P., Pesti, G., Bakalli, R. and Edwards, H. (2006). The effect of feeding calcium-and phosphorus deficient diets to broiler chickens during the starting and growing-finishing phases on carcass quality. *Poultry Science*. 85: 1939-1946.
- Duncan, D. B. (1955). Multiple Range and Multiple F Tests. *Biometrics*, 11, pp.1-42.

- Eeckhout, W., and Peape, M. D. (1994). Phytata-und Phytata pin Futtermitein. *Animal Feed Science and Technology*. 47: 19-29.
- Greiner, R. and Konietzny, U. (2011). Phytases: Biochemistry, enzymology and characteristics relevant to animal feed use. In: *Enzymes in Farm Animal Nutrition* (2nd ed.). Eds. Bedford, M. & Partridge, G., CABI publishing, Wallingford, Oxfordshire, UK. pp. 96-128.
- Guggenbuhl, P., Torrallardona, D., Cechova, I., SimoesNunes, C., Wache, Y., Fru, F. and Broz, J. (2012). The efficacy of a novel microbial 6-phytase expressed in *aspergillusoryzae* on the performance and phosphorus utilization in swine. *Journal of Animal Science Advertisement*. 2: 438-452.
- Hastings, W.H. (1946). Enzyme supplements to poultry feeds. *Poultry Science*. 25:584-586.
- Hartig, T. (1855). Uber das klebermehl. *Botanische Zeitung*. 13: 881-885. (cited by Reddy *et al.*, 1982).
- Hídvégi, M. and Lasztity, R. (2002). Phytic acid content of cereals and legumes and interaction with proteins. *Periodica Polytechnica Ser. Chem. Eng.* 46: 59-64.
- Howlider, N. and Rose, K. (1989). Evaluation of microbial phytase in broiler diets. *Poultry Science*. 83, 962–970.
- Islam, F.A., Simon, O., Milksch, G. and Manner, K. (2010). Comparative studies of the *in vitro* properties of phytases from various microbial origins. *Arch. Anim. Nutr.* 53, 353–373.
- Jaap, P. H., Ravindran, V. and Pittolo, P. H. (1950). An evaluation of microbial phytase in sorghum-based broiler diets. *The Proceedings of the Australian Poultry Science Symposium*, 11:97–100.
- Jalal M., Scheideler S. and Pierson E. (2001). Strain response of laying hens to varying dietary energy levels with and without Avizyme supplementation. *J. Appl. Poult. Res.* 16:289–295.
- Jeroch, H. (1993). *Futtermittellekinde*. Gustav Fischer Verlag, Jena-Stuttgart, 307–322.

- Johnson, L. & Tate, M.(1969). Structure of "phytic acids". *Can. J. Chem.* 47: 63-73.
- Jordao, F. J., Vilar, D. S., Silva, J. H., Eernandes, A. C.; Telxeira, E. N. M., Ribeiro, M. L. G. and Costa, F. G. P. 2006. Níveis de P disponível e de fitase na ração de frangos de 22 a 42 dias. *Revista Brasileira de Ciencia Avicola.* 8:107
- Kerr, B., Weber, T., Miller, P. and Southern, L. (2010). Effect of phytase on apparent total tract digestibility of phosphorus in corn-soybean meal diets fed to finishing pigs. *J. Anim. Sci.* 88: 238-247.
- Khawaja, Z. (2003). Utilization of citrus pulp in broiler rations. Master's thesis. Faculty of Graduate Studies, An Najah National University.
- Kies, A. K., Vanhemert, K. A. F. and Sauer, W.C. (2001). Effect of phytase on protein and amino acid digestibility and energy utilization. *World's Poult. Sci. J.* 57: 109-124.
- Knuckles, K. and Betschart, T. (1987). The analysis for inositol phosphate forms in feed ingredients. *J. Sci. Food Agric.* 76, 1-9.
- Knuckles, B. (1989). Effect of Myo inositol phosphateesters on in vitro and in vivo digestionof protein. *Journal Food Science.* 54(5): 4-9.
- Kornegay Z., Yi, E.T. and Ravindran, V. (1996). Phytase and phosphorus availability. *Poultry Science.* 75: 240-249.
- Kundu B., Biswas P., Rajendran D. and Dandopath S.K. (2000). Effect of supplementation of phytase enzyme and gradual re-placement of dicalcium phosphate on the performance of commercial broiler chicken. PP: 119-128 in Proc. III Biennial conference of Animal Nutrition Association, Hisar, India.
- Kwon, K., Chag, J.I., Ju, J.C., Yoo, M.I., Sohn, K.S. and Choi, Y.W. 1999. Effects of microbial phytase on performance of broiler chicks fed corn-soy based diets. *Korean journal of Animal Science.* 41:519-526.
- Lan, G. Q., Abdullah, N., Jalaludin, S., and Ho, Y. W. (2002). Efficacy of supplementation of a exogenous phytase in low protein diet of broiler. *Poultry Science.* 81(10): 1522-1532.

- Lott, J., Ockenden, I., Raboy, V. and Batten, G. D. (2000). Phytic acid and phosphorus in crop seeds and fruits: A global estimate. *Seed Sci. Res.* 10: 11-34.
- Lowe, J.T., Steenbock, H. and Keiger, C.H., 1939. Cereals and rickets. IX. The availability of phytin-P to the chick. *Poultry Science.* 18, 40–44.
- Maenz, D. (2001). Enzymatic characteristics of phytases as they relate to their use in animal feeds. In: *Enzymes in Farm Animal Nutrition*. Eds Bedford, M. & Partridge, G., CABI publishing, Wallingford, Oxfordshire, UK, pp: 61-84.
- McNally, S. and Spicknall, I. (1949). Phytate and phytase: consequences for protein utilization. *Nutr. Res. Rev.* 13:255–278.
- Miles, R.D. and Nelson, T. S. (1974). The effect of enzymatic hydrolysis of phytate on the available energy content of feed ingredients for chickens and rats. *Poultry Science.* 53: 1714-1717
- Mondal, M. K., Panda, S. and Biswas, P. (2007). Effect of microbial phytase in soybean meal based broiler diets containing low phosphorous. *Internal Journal of Poultry Science.* 6 (3): 201-206.
- Moshad, S. H. (2001). Advances in the development and application of feed enzymes. *Australian Poultry Science. Symp.* 7: 109–115
- Morkunas, M., Kbtitskas, G. and Kublitsken, V. (1993). An enzyme premix. *Poultry Abstract.* 19: 43.
- Mullaney, E. J., Daly, C. B. and Ullah, A. H. (2000). "Advances in phytase research". *Advance Applied Microbiology.* **47**: 157–199.
- Mullaney, E. J. and Ullah, A. H. (2003). "The term phytase comprises several different classes of enzymes". *Biochem Biophys Res Commun.* **312** (1): 179–184.
- Naher, B. (2000). Utilization of parboiled rice polish based diet with supplementation of carbohydrate and phytase in growing ducklings. M.S. Thesis, Department of Poultry Science, Bangladesh Agricultural University, Mymensingh.

- Namkung, H. and Leeson, S. (1999). Effect of phytase enzyme on dietary nitrogen-corrected apparent metabolisable energy *and* the ileal digestibility of nitrogen *and* amino acids. *Poultry Science*. **78**: 1317–1319.
- National Research Council, (1994) Nutrient requirements of Poultry. 9<sup>th</sup> rev. ed. National Academy Press, Washington, DC.
- Nadeem, M. A., Anjum, M. I., Khan, A. G. and Azim, A. 2005. Effect of dietary supplementation of non- starch polysaccharide degrading enzymes on growth performance of broiler chicks. *Pakistan Veterinary Journal*. 25(4): 183–187.
- Orban, D.J., Martin, E., Preez, J. J., Bongarts, M., Betts, M., Sudaman, A. and Thomson, E. (1999). The beneficial effects of a microbial phytase in diets of broiler chicken, duckling and turkey. *J. Anim. Physiol. Animal Nutrition*. 69: 278–283.
- Oloffs, K., Cossa, J. and Heroch, H. (2000). Phosphorus utilization different vegetable feedstuffs by laying hens//*Arch. Geflugelk*. 64(1) 24-27.
- Piao et al.; (1998). Phosphorus requirements of broilers and the impact of exogenous phytases. *Ark. Nutr. Conf*.
- Pizzolante, C., Oduguwa, O., Acamovic, T. and Bedford, M.R. (2008). Effects of dietary phytase on performance and nutrient metabolism in chickens. *British Poultry Science*. 49: 144-154.
- Plumstead P.W., Leytem A.B. and Maguire R.O. (2008). Interaction of calcium and phytate in broiler diets. Effects on apparent prececal digestibility and retention of phosphorus. *Poultry Science*. 87: 449-458.
- Pillai, A. R., Venegowda, G. and Aravin, B.I.R. (1995). Influence of enzyme supplementation on the performance of broiler fed tannin-rich diets. *International Journal Poultry Science*., 30: 248-250.
- Pillai, P. B., Conner- Dennie, T. O., Owens C. M. and Emmert, J. L. (2006). Efficacy of an E. Coli Phytase in broiler fed adequate or reduced phosphorus diets and its effects on carcass characteristics. *Poultry Science*. 85: 1200- 1211.



- Qian, H., Vett, H. P., Kornegay, E. T., Ravindran, V. and Denbow, D. M. (1996). Effects of supplemental phytase and phosphorus on histological and other tibial bone characteristics and performance of broilers fed semi-purified diets. *Poultry Science*. 75 (5): 618-626.
- Rahman, A., Saima., Pasha, T. N., Younus, M., Abbasi, Y. and Allah, Y. (2009). Effects of supplemental microbial phytase on performance of broiler chickens. *Agriculture Science in Finland* 3:457-466.
- Rabayaa, E. (2003). Utilization of olive cake in broilers ration. Master's thesis. Faculty of Graduate Studies. An Najah National University.
- Rama Rao, A., Slominski, B. A., Jia, W., Nyachoti, C. M. and Wittenberg, K. M. (1999). Apparent metabolizable energy (AMEn) content and standardized ileal amino acids digestibility of wheat, wheat-corn and corn distillers dried grains with soluble (DDGS) for broilers. *Poultry Science*. 89: 556.
- Rama Rao, S., Raju, M., Reddy, M. and Pavani, P. (2001). Interaction between dietary calcium and non-phytate phosphorus levels on growth, bone mineralization and mineral excretion in commercial broilers. *Animal Feed Science Technology*. 131: 135-150.
- Ravindran, V., Cabahug, S., Ravindran, G. and Bryden, W. (1999). Influence of microbial phytase on apparent ileal amino acid digestibility of feedstuffs for broilers. *Poultry Science*. 78: 699-706.
- Ravindran, V., Cabahug, S., Ravindran, G., Selle, P. and Bryden, W. (2000). Response of broiler chickens to microbial phytase supplementation as influenced by dietary phytic acid and nonphytate phosphorous levels. II. Effects on apparent metabolisable energy, nutrient digestibility and nutrient retention. *Broiler Poultry Science*. 41: 193-200.
- Ravindran, V., Selle, P., Ravindran, G., Morel, P., Kies, A., and Bryden, W. (2001). Microbial phytase improves performance, apparent metabolizable energy, and ileal amino acid digestibility of broilers fed a lysine-deficient diet. *Poultry Science*. 80: 338-344.

- Reddy, N. and Sathe, S. D. (1982). Phytates in legumes and cereals. *Advance Food Reseach.* 28: 1-75.
- Ren, Q. M., Zahn, H., Song, G. and Zhan, H.Y. (1999). A study on nutritional effects of subnormal dietary phosphorus supplementation of microbial phytase on broilers. *Acta Agriculture Univesitatis Jiangxiensis*, 21: 575-580
- Richered, A.J., Acamovic, T. and Bedford, M. R. (1994). The effect of phytase and phytic acid on endogenous losses from broiler chickens. *British Poultry Science.* 45: 101-108
- Rosen, M. K. (2010). Enzymes in poultry diets in : *Enzymes in Animal Nutrition. Proceedings 1<sup>st</sup> Symposium, Switzerland. Oct 13-16.* pp: 61-63.
- Rojas, S. and Scott, M. (1969). Factors affecting the nutritive value of cottonseed meal as a protein source in chick diets. *Poultry Science.* 48: 819-835.
- Sauvant, D., Prerez, J. M. and Tran, G. (2004). *Tables of composition & nutritional value of feed materials: Pigs, poultry, cattle, sheep, goats, rabbits, horses and fish (2nd ed.) Academic Pub. Wageningen.*
- Selle, P., Ravindran, V., Pittolo, P. and Bryden, W. (1999). An evaluation of microbial phytase in sorghum-based broiler diets. *Proc. Aust. Poult. Sci. Symp.* 11 . pp. 97-100.
- Selle, P., Ravindran, V., Caldwell, R. and Bryden, W. (2001). Phytate and phytase: Consequences for protein utilization. *Nutr. Res. Rev.* 13: 255-278.
- Selle, P. H. and Ravindran, V. (2005). Microbial phytase in poultry nutrition. *Animal Feed Science and Technol.* 135: 1-41.
- Selle, P. H., Walker, A. R. and Bryden, W. L. (2006a). Total *and* phytate-phosphorus contents *and* phytase activity of Australian sourced feed ingredients for pigs *and* poultry. *Aust. J. Expt. Agric.* **45**: 475–479.
- Selle, P. H., Ravindran, V., Bryden, W. L. and Scott, T. (2006b). “Influence of dietary phytate and exogenous phytase on amino acid digestibility in poultry, a review,” *The J. Poult. Sci.* 43(2): 89–103.

- Selle, P. H. and Ravindran, V. 2007. Microbial phytase in poultry nutrition. *Animal Feed Science and Technology*. 135: 1-41.
- Sebastian, S., Touchburn, S. P., Chavez, E. R. and Lague, P. (1996). Efficacy of supplemental microbial phytase at different dietary calcium levels on growth performance and mineral utilization of broiler chickens. *Poultry Science*., 75: 1516-1523.
- Sebastian, S., Touchburn, S. P., Chavez, E. R. and Lague, P. C. (1997). Apparent digestibility of protein and amino acids in broiler chickens fed a corn-soybean diet supplemented with microbial phytase. *Poultry Science* 76:1760-1769.
- Sohail, S.S. and Roland, D.A. 1999. Influence of supplemental phytase on performance of broilers four to six weeks of age. *Poultry Science*. 78: 550-555.
- Scott, T. A., Swift, M. L. and Bedford, M. R. (1997). The influence of feed milling, enzyme supplementation and nutrient regimen on broiler chick performance. *J. Appl. Poult. Res.* 6: 391-398.
- Shelton, J., Southern, L., Gaston, L. and Foster, A. (2004). Evaluation of the nutrient matrix values for phytase in broilers. *J. Appl. Poultry Res.* 13: 213-221.
- Shirley, R. and Edwards, H. M. Jr., 2003. Graded levels of phytase past industry standards improves broiler performance. *Poultry Science*. 82: 671-680.
- Simons, P. C. M., Versteegh, H. A. J., Jongbloed, A. W., Kemme, P. A., Slump, P., Bos, K. D., Wolters, M. G. E., Beudeker, R. F. and Verschoor, G. J. (1990). Improvement of phosphorus availability by microbial phytase in broilers and pigs. *British Journal for Nutrition*. 64:525-540.
- Simon, O. and Igbasan, F. (2002). In vitro properties of phytases from various microbial origins. *Internal Journal of Food Science and Technology*. 37: 813-822.
- Singh P.K. and Khatta V.K. (2002). Phytase supplementation for economic and eco-friendly broiler production. *J. Eco. Physiol.* 5, 117-121.

- Singh P.K. and Khatta V.K. (2003a). Effect of phytase supplementation on the performance of broiler chickens fed wheat based diets. *Industrial Journal for Animal Nutrition*. **20**, 57-62.
- Singh P.K. and Khatta V.K. (2003b). Effect of phytase supplementation on the broiler production. *Poult. Plan*. **4**, 13-14.
- Singh P.K. and Khatta V.K. (2004). Economics of broilers raised on phytase supplemented diets. *Industrial Journal Animal Reseach*. **10**, 121-124.
- Singh, P. (2008). Significance of phytic acid and supplemental phytase in chicken nutrition: A review. *World's Poultry Science*. J. 64: 553-580.
- Vinil S.P., Kadirvelan C., Thirumalai S., Valli S. and Chandrabose B. (2000). The role of phytase and other enzymes in augmenting the bioavailability of nutrients in broilers. Pp. 97 in Proc. xx Annual Conference of Indian Poultry Science Association, Chennai, India.
- Van Der Klis, J. D., Versteegh, H. A. J. and Simons, P.C.M. (1997). The efficacy of phytase in corn-soybean meal-based diets for laying hens. *Poultry Science*.76:1535-1542.
- Warden, D. and Schaible, k. (1962). Effect of dietary phosphorus, phytase, and 25-hydroxycholecalciferol on broiler chicken bone mineralization. *Poultry Science*. 85:1200-1211.
- Wilson, W., Chicco, C., Meschy, F. and Requena, F. (1999). Phytic phosphorus and phytase activity in chickens. *Interciencia*, 30: 24-28.
- Watson, B. C., Matthews, J. O., Southern, L. L. and Shelton, J. L. 2006. "The effects of phytase on growth performance and intestinal transit time of broilers fed nutritionally adequate diets and diets deficient in calcium and phosphorus," *Poultry Science*. 85(3): 493-497.
- Zhang, Z., Kornegay, E., Radcliffe, J., Wilson, J. and Veit, H. (2000). Comparison of phytase from genetically engineered aspergillus and canola in weanling pig diets. *Journal of Animal Science*. 78: 2868-2878.
- Zyla, K. (1992). Mould phytases and their application in the food industry // *World Journal of Microbiology and Biotechnology*. 8: 467-472.